

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

Primary IV Fluid and Infusion Rate (ml/hr.)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
LINDSEY LAMBERT	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr. and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Oxytocic Hormone (synthetic substitute)	Induce labor	1 mU/min Max 20U/min	Y N	D5W NS	Fetal trauma/death Anaphylactic reaction edema	1. Report sudden, severe headache immediately to HCP 2. Start continuous FHM 3. If epidural given + oxytocin, monitor for hypertensive crisis 4. Monitor I & O- water intoxication
Magnesium Sulfate	Anticonvulsant electrolyte	Inhibit premature labor- tocolytic	4g in 250ml	Y N	D5W	Sweating Extreme thirst Depressed reflexes	1. Observe constantly when IV- BP/pulse 2. Monitor plasma magnesium levels 3. Magnesium toxicity- cathartic effect/ thirst/ warm/sedation/depressed DVT 4. Have calcium gluconate readily available
Meperidine	CNS agent- Opiate	Control moderate-severe pain	50-100mg Q1-3h	Y N	NS Sterile water D5W	Allergic reaction Profuse perspiration dizziness	1. Give smallest effective dose first 2. Assess pt need for PRN medication 3. Monitor vitals closely (HR, BP) 4. Repeated use can lead to tolerance
Promethazine	Antiemetic	Control nausea and vomiting	12.5-25mg Q4-6hr PRN	Y N	NS	Deep sleep Drowsiness Blurred vision	1. Do not mix with alcohol or other CNS depressants 2. Supervise ambulation- sedation/dizziness 3. Avoid prolonged sun exposure 4. Pt may develop involuntary movements
Calcium Gluconate	F&E and water	Hypocalcemia Hypermagnese	1.5-3 g IV	Y	Undiluted	Tingling sensation	1. Monitor ECG during IV administration 2. Observe IV site closely

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

	balance agent	mia Hyperkalemia	2-5 min	N	NS	Constipation Hypotension	3. Lab tests: Calcium, phosphorus, magnesium 4. Monitor BP- Hypotension
Phytonadione	Vitamin K	Prophylaxis & therapy for neonatal hemorrhagic disease	0/5-1mg IM immediately after delivery	Y N	n/a	Pain at injection site Skin rash Hypersensitivity Hyperbilirubinemia	1. NB are unable to produce vit K needed for certain coagulation factors 2. Provide comfort during and after- kangaroo hold/swaddle 3. Admin in vastus lateralis 4. Must get consent
Erythromycin Ophthalmic Ointment	Anti-infective	Prophylaxis of ophthalmia neonatorum	0.51cm ribbon in eyes after birth	Y N	n/a	Burning Tenderness dryness	1. Use new tube for each neonate 2. Culture site of infection before therapy if indicated 3. Pull lower eyelid down and admin inner to outer corner 4. avoid touching eye
Engerix B	Vaccine	Hepatitis B active immunity	0.5 ml at birth IM	Y N	n/a	Mild tenderness at site Local inflammatory reaction	1. Give vaccine information sheet 2. Must obtain consent 3. Admin in vastus lateralis 4. Document time, date, location, consent, education, and lot number
Hepatitis B Immune Globulin	Vaccine	Passive immunity to neonates born to hep b + moms	0.5ml at birth IM	Y N	n/a	Muscle stiffness Tenderness Rash	1. Give at same time at hep B vaccination 2. Admin in vastus lateralis 3. Must obtain consent 4. Document time, date, location, consent, education, and lot number