

Student Name: ASHLEE GUZMAN

Unit: \_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: 10/11/21

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: N/A

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/Hypotonic/Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push.	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Pitocin	labor induction, hemorrhage	0.5-1 mu per min IV DISKONIN	<input checked="" type="checkbox"/> Y <input type="checkbox"/> N	Dilute to 10 mu/ml by adding 10u to 1000ml of D5W, LR, NS	neonatal seizure, jaundice to neonatal, low Apgar score, fetal hypoxia, tachycardia	1. ALWAYS monitor fetal heart rate before/during/after. 2. Have sure cervical assessment is done. 3. If there is fetal distress, stop oxytocin. 4. Do Bishop scoring to check cerv. dilation.
Magnesium Sulfate	Anti-hypertensive electrolyte	eclampsia, preeclampsia, arrhythmia,	1g/100 ml Q4 hrs.	<input checked="" type="checkbox"/> Y <input type="checkbox"/> N	infuse over 2-4hr rate not to exceed 125mg/hr	pulmonary edema, hypotension, flushing, drowsiness, circulatory collapse	1. Prior to admin. check serum magnesium level. 2. Have patients report breathing difficulties. 3. Monitor pt. for magnesium toxicity. report loss of deep tendon reflexes. 4. Have pt. report respiratory depression.
Meperidine	Demerol	OPioids ANALGESICS	50mg TAB. PRN	<input checked="" type="checkbox"/> Y <input type="checkbox"/> N		agitation, angina, bradycardia, cardiac arrest, coma, dry mouth	1. Monitor for respiratory depression. 2. Be ready to resuscitate baby after labor. 3. Teach patient prolong use can cause baby to opioid withdrawal. 4. to report any dizziness.
Promethazine	Prenergan	Anti-histamine	12.5 mg TAB. twice daily	<input checked="" type="checkbox"/> Y <input type="checkbox"/> N		sedation, confusion, blurred vision, hallucinations, dystonia, disorientation	1. Teach pt. caution if driving going to drive car impair. 2. Teach pt. to report symptoms of drowsiness. 3. Teach pt. to report any fever. 4. Teach pt. to report dizziness.
Calcium Gluconate	gluconate	antidotes	50mg TAB. Q8-12hr	<input checked="" type="checkbox"/> Y <input type="checkbox"/> N		bradycardia, hypotension, muscle, constipation, diarrhea, N/V	1. Teach pt. that constipation may occur. 2. Avoid if pt has hypercalcemia. 3. Monitor pt. for cardiac arrest. 4. Use with caution in pts. w/ severe hyperphosphatemia.

Student Name: ASHLEE GUZMAN

Unit: \_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: 10/11/21

Allergies: N/A

Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications				
	Isotonic/Hypotonic/Hypertonic							
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push.	IVPB - List ml/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	MEMO-STATICS	VITAMIN K	100 mcg Daily +tbl.	<input checked="" type="checkbox"/> Y			dyspnea, cyanosis, flushing, hypotension, pruritus	<ol style="list-style-type: none"> <li>Never admin. a fast infusion.</li> <li>TEACH pt. to report severe reactions.</li> <li>Proper dilution has to be administered.</li> <li>Monitor for respiratory arrest.</li> </ol>
Erythromycin Ophthalmic Ointment	MACRO-VIDES	ANTI-BIOTIC	0.500 1hr after labor	<input checked="" type="checkbox"/> Y			hypersensitivity reactions, redness, ocular reactions	<ol style="list-style-type: none"> <li>TEACH pt. it has to be put on within 1hr of labor.</li> <li>Don't put if ocular irritation occurs, let tip of applicator touch the eye.</li> <li>Always avoid contamination</li> </ol>
Engerix B	HEPLISAV-B	INACTIVATED VACCINE HEPATITIS B	1mL IM 3-dose series	<input checked="" type="checkbox"/> Y			Pain, fatigue, pyrritis, erythema, burning	<ol style="list-style-type: none"> <li>Never give IV or intrac dermal.</li> <li>Always admin in IM deltoid muscle it's an infant w/ admin anterolateral thigh.</li> <li>TEACH pts it's a 3-dose series.</li> </ol>
Hepatitis B Immune Globulin	IMMUNE GLOBULINE	ANTIBODIES	512 units/mL IM	<input checked="" type="checkbox"/> Y			headache, erythema, nausea, vomiting, malaise	<ol style="list-style-type: none"> <li>NEED to separate live vaccines by 3months. decrease.</li> <li>Monitor WBC if may decrease.</li> <li>USE caution w/ pts who have bleeding disorders.</li> </ol>