

References

- Lester, B. M., & Twomey, J. E. (2008). Treatment of substance abuse during pregnancy. *Women's Health*, 4(1), 67–77. <https://doi.org/10.2217/17455057.4.1.67>
- Murray, S. S., McKinney, E. S., Holub, K. S., & Jones, R. (2019). Saunders. *Foundations of maternal newborn & women's health nursing*.
- Pinto, S. M., Dodd, S., Walkinshaw, S. A., Siney, C., Kakkar, P., & Mousa, H. A. (2010). Substance abuse during pregnancy: Effect on pregnancy outcomes. *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 150(2), 137-141. <https://doi.org/10.1016/j.ejogrb.2010.02.026>
- Roloff, K. A. (2015, September 19). ClinMed International Library. *Methamphetamine use increases chances of preeclampsia*, 1(3). <https://doi.org/10.23937/2469-5793/1510011>
- Shankaran, S., Das, A., Bauer, C. R., Bada, H. S., Lester, B. M., Wright, L. L., Higgins, R. D., & Pool, W. K. (2011). Neurotoxicology and teratology. *Prenatal cocaine exposure and small-for-gestational-age status: Effects on growth at 6 years of age*, 33(5), 575-581. <https://doi.org/10.1016/j.ntt.2011.04.003>