

**Maternal Medication Worksheet – Current Medications & PRN for Last 24 Hours**

**Allergies:** \_\_\_\_\_

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push.  IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Uterine smooth muscle stimulant	Stimulates uterine contractions	0.5-1 milliunit/min  1-2milliunits/min q30-60 mins until desired contraction pattern is established	Y  N	IVPB-- dilute in 1,000 mL of NS or LR  Rate of admin as ordered by physician	<b>Black box:</b> Not to be given for elective labor induction, but can be used when there is a clear medical indication for induction  Hypertonicity may occur with tearing of uterus, increased bleeding, abruptio placentae, cervical/vaginal lacerations	1. Monitor B/P, pulse, respirations, fetal heart rate, intrauterine pressure, contraction q15 min  2. Notify physician of contractions that last longer than 1 minute, occur more frequently than every 2 minutes, or stop  3. Keep pt informed of labor progress  4. Assess baselines for vital signs, B/P, fetal heart rate and determine frequency, duration, and strength of contractions
Magnesium Sulfate	Anticonvulsant	Prevention and treatment of seizures in severe preeclampsia or eclampsia	4-6 g over 20-30 min IV	Y  N	dilute to maximum concentration of 20% for IV infusion	N/V  diarrhea  large drops in BP  breathing	1. Monitor for deep tendon reflex loss  2. Monitor respiratory status  3. Monitor B/P  4. Teach not to take with hypertensive medications

Student Name: \_\_\_\_\_

Unit: \_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: \_\_\_\_\_

**Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

						difficulty	
Meperidine	Opioid	Relieve moderate to severe pain	PO 50mg/5mL 50 mg, 100mg	Y N	25mg/ml 100mg/ml	constipation, depression of cough reflex, sedation	1. Suddenly stopping may cause withdrawal 2. Teach pt about risk for addiction and abuse 3. Avoid if renal impairment 4. Teach antidote is naloxone
Promethazine	Phenothiazine	treatment or prevention of N/V	12.5-25 mg PO, PR, IM q 4-6 hours	Y N	100mg/mL IV 500mg PO	extreme drowsiness, neuroleptic malignant syndrome	1. Monitor platelets and wbc 2. Assess breathing frequently 3. teach EPS 4. teach interaction with other medications
Calcium Gluconate	Antidotes	treat hypocalcemia	1000/day orally q 8-12 hours	Y N	100mg/mL IV 500mg PO	dry mouth, headache, hot flashes, nausea	1. teach not to start, stop or change dosage before checking with provider 2. caution if breastfeeding 3. take with food 4. report bradycardia

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	hemostatics	vitamin	1mg/min IV	Y N	dilute in NS 50 mL over 20 min	CNS depression, respiratory arrest, cardiac arrest	1. monitor respiratory status frequently 2. be informed about cardiac issues 3. report difficulty breathing 4. report rash
Erythromycin Ophthalmic Ointment	macrolide antibiotic	prevent eye infections in newborns	ointment	Y N		redness, itching, stinging, burning of the eye	1. Used one time soon after delivery to prevent eye infection 2. follow prescription label carefully 3. clean hands thoroughly before applying 4. wipe excess ointment from eyelids

Student Name: Airykah Molina

Unit: \_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: 10/18/21

**Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**

**Allergies:** \_\_\_\_\_

Engerix B	vaccine	prevent hep B infection	10mcg/0.5MI	Y N	10mcg/0.5MI	allergic reaction, irritability, drowsiness, high temp	1. give first injection at birth 2. injection given into thigh 3. teach may cause bruising 4. teach to report signs of jaundice
Hepatitis B Immune Globulin	immune globulin	protect babies born to mother who test positive for hep B	IM	Y N	0.5 mL	fever, nausea, diarrhea, hives	1. monitor for 15 minutes after given 2. finish all doses of hep b vaccine 3. follow up blood tests 4. give within 7 days
				Y N			1. 2. 3. 4.