

Student Name: Michael Aranda

Maternal Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: N/A

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin (Pitocin)	Exogenous hormone	To induce or stimulate labor	0,5 – 1 mu per min IV Q 15-60 min	Y N	Dilute to 10 mu/ml by adding 10 u to 1000ml of D5W, NS, or LR	Seizure, subarachnoid hemorrhage, coma, arrhythmia, abruptio placentae	1. CAUTION: may cause anti-diuretic effect and risk of severe water intoxication, seizures or death. 2. ALERT: all patients receiving oxytocin must be under continuous observation by trained professional. 3. REPORT: blurred vision, itching, difficulty speaking, and nausea 4. ALERT: discontinue oxytocin immediately if uterine hyperactivity or fetal distress occurs.
Magnesium Sulfate	Magnesium salts	laxative	1g/ 100ml Q 4 hours	Y N	Infuse over 2-4 hours. Rate not to exceed 125 mg/hour	Abdominal cramping, diarrhea, nausea, fluid and electrolyte disturbances	1. CAUTION: may alter fluid and electrolyte levels with prolonged use 2. ALERT: watch for magnesium toxicity 3. Report: slurred speech, sweating, chest pain, and difficulty breathing. 4. Ensure calcium gluconate is readily available.
Meperidine (Demerol)	Opioid	Opioid analgesic	50 mg tab PRN	Y N		Bradycardia, cardiac arrest, shock, respiratory arrest, anaphylaxis	1. CAUTION: may increase amylase, lipase, and CK levels. 2. Call for help when getting out of bed or walking. 3. ALERT: don't stop drug abruptly; withdrawal slowly 4. Monitor pt. for adrenal insufficiency: N/

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

							V, loss of appetite, fatigue, & weakness
Promethazine (Phenergan)	Phenothiazines	Antiemetic	12.5 mg tab twice daily	Y N		Leukopenia, agranulocytosis, thrombocytopenia, respiratory depression, and apnea	<ol style="list-style-type: none"> 1. CAUTION: may increase Hb level and hematocrit and blood glucose level. 2. Take drug with milk or food 3. Report involuntary muscle movements 4. May decrease WBC, platelet and granulocyte counts.
Calcium Gluconate	Calcium salts	Calcium supplement	50 mg tab Q 8-12 hour	Y N		Tingling sensation, hemorrhage, cardiac arrest, bradycardia, vasodilation	<ol style="list-style-type: none"> 1. CAUTION: may increase calcium levels 2. Take with a full glass of water 3. Do not eat rhubarb, spinach, bran or whole grain cereal (interferes w/ absorption) 4. Report: dry mouth, thirst, or polyuria

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione (vitamin k)	Vitamins Hemostatic	Prevent bleeding by helping to clot	100 mcg daily tab	Y N		Taste changes, dizziness, tachycardia, sweating, SOB, and flushing.	<ol style="list-style-type: none"> 2. ALERT: immediate coagulant effect should not be expected after administration. 2. WILL NOT counteract the anticoagulant action of heparin 3. Protect from light or it will degrade rapidly 4. Dosage should be kept as low as possible
Erythromycin Ophthalmic Ointment	Macrolides	Antibiotics	0.5% 1 hour after	Y N		Minor ocular irritation, burning,	<ol style="list-style-type: none"> 1. ALERT: prolonged use may result in fungal or bacterial superinfection. 2. Gently massage eyelids for 1 minute to spread ointment.

Student Name: Michael Aranda

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: N/A

			labor			dryness, oily skin, peeling	3. Don't use for infection unless causative organism has been identified. 4. To prevent ophthalmia neonatorum, apply ointment no later than 1 hour after birth.
Engerix B (hep B adult vaccine)	Hep B vaccine	Inactivated vaccine (Hep B)	1mL IM 3-dose series	Y N		Fever, HA, tiredness, sore throat, nausea, diarrhea, loss of appetite, and dizziness	1. CAUTION: avoid if allergic to yeast 2.ALERT: will not protect against HEP B if already infected 3.Teach: will not protect against hep A, C, or E, or other viruses that attack the liver 4.Report: light headedness or feeling like you are going to pass out and involuntary muscle movements
Hepatitis B Immune Globulin	Immune serum	Prophylaxis drug	312 units/mL IM	Y N		Chills, fever, HA, malaise, hypotension, low back pain	1. CAUTION: may increase creatinine level and LFT values. 2. Report tenderness at injection site. 3. Keep epinephrine and antihistamine available 4. CAUTION: can interfere with certain types of blood glucose monitoring systems and cause falsely elevated readings.
				Y N			1. 2. 3. 4.