

IM 2 Student Professional Skills Video Grading Rubric

Student Name: __Mikah Barnes__

Date: __11/07/21__

Universal Competency	Critical Elements	M	U
Safety (physical and emotional)	Introduce self	X	
	Identify patient	X	
	AIDET		X
	Allergy	X	
	4 P's	X	
	Medication Administration		
	Medication, dosage, route, reason		X
Critical Thinking	Assessment: Focused		
	Perform according to professional nursing intervention instruction	X	
	Procedure		
	Assess, Plan, Implement, Evaluate (APIE)	X	
	Correct equipment	X	
Standard Precaution	Asepsis		
	Hand hygiene	X	
	Don gloves if indicated		
	Clean equipment (stethoscope, pulse oximeter, etc.)		
	Procedures Medication preparation/delivery		
Communication	Teach patient		
	Medication Procedure		X
Documentation	Scan patient		X
	Scan medication		
	Assessment findings		
	Procedure		
	Save documentation		
Human Caring and Relationship	Respect, active engagement, authenticity, empathy	X	
Professional Role Performance	Appearance, preparation, behaviors, resource management	X	
Instructor comments:			

Created by IM2 Faculty, Covenant School of Nursing, Feb. 2021

Unmet Comments:

- AIDET: I did not address the D of aidet so I did not say how long I would be in the room, which I would let my patient know I'd be in the room for approximately 20 minutes performing my assessments and giving medications.
- Documentation: I did not state that I would have documented the findings associated with the nasal cannula and skin breakdown, and for the skin on both the chest for the

cardiac focused assessment, and the abdominal area for the GI focused assessment. However I did scan the wristband and medications and saved my documentation.

- Medication Administration: I did not tell patient the reason for why I was giving the metoprolol medication, but I would have told my pt. it is a med for her heart rate and her history of bradycardia. I also did not educate my patient to eat within 15 min of administering the insulin but I would have done so, the reason being I don't want my pt. to bottom out when the medication is delivered. Also for insulin I would have explained to the pt. that the injection would be 2in away from the umbilicus, or the middle of the abdomen and I would look out for any lipodystrophy. When giving the medications through the tube I would have placed a towel down on the pts. stomach to ensure they did not get wet during the administration.