

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
Click here to enter text.	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>	Click here to enter text.	Click here to enter text.	Click here to enter text.

Student Name: Rushondra S		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	Allergies: Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
(Zofran) Ondansetron	Antiemetics	To prevent n/v from highly emetogenic To prevent n/v from moderately emetogenic chemotherapy To prevent postoperative n/v N/V during pregnancy	PO, IM, IV	Choose an item. Click here to enter text.	Dilute drug in 50 ml of d5w, infuse over 15 minutes	CNS: dizziness, fatigue, headache, malaise CV: arrhythmia, chest pain GI: constipation, diarrhea GU: gynecologic Respiratory: hypoxia Skin: pruritus	1. An ekg may be needed to monitor heart rate and rhythm . 2. Report difficulty breathing after drug administration 3. Contact provider immediately if he experiences s/s of abnormal heart rate. 4. Place medication on tongue allow to dissolve, then swallow with saliva
(Cuvposa) Glycopyrrolate	Anticholinergics	used to reduce saliva and drooling in children	IV, IM	Choose an item. Click here to enter text.	0.1mg iv repeat 2 to 3 minutes		1. Take an empty stomach. 2. Take 1 to 2 hours after a meal. 3. Measure liquid medicine carefully 4. Use a medicine dose - measuring device

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

(Anectine) Succinylcholine	Neuromuscular blocking agents	Relax skeletal muscle for surgery and orthopedic manipulations, to assist with mechanical ventilation	IV	Choose an item. Click here to enter text.	0.6mg /kg give over 10 to 30 seconds	CV: arrhythmias, bradycardia , cardiac arrest EENT: increased intraocular pressure GI: excessive salivation Metabolic: hyperkalemia Respiratory: apnea, bronchoconstriction, prolonged respiratory depression.	<ol style="list-style-type: none"> 1. Explain all events and procedures to pt because he can still hear. 2. Reassure patient that postoperative stiffness is normal and will soon subside.
(Diprivan) Propofol	Anesthetics	To maintain anesthesia, monitored anesthesia, to sedate intubated intensive care unit ICU patients	IV	Choose an item. Click here to enter text.	5mcg/kg/min for 5 min, increments of 5 to 10mcg/kg/min	CNS: dystonic or choreiform movement CV: bradycardia Metabolic: hyperlipidemia Skin: rash, pruritus Other: burning or stinging at injection site	<ol style="list-style-type: none"> 1. Do not perform activities requiring mental alertness 2. Urine may be turned green 3. Abnormal dreams or anesthesia awareness may occur 4. Stop drug gradually

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

<p>(Lipitor) Atorvastatin</p>	<p>Antilipemics</p>	<p>Help reduce the risk of of nonfatal MI, fatal and non-fatal strokes, angina, heart failure, and revascularization procedures</p>	<p>PO</p>	<p>Choose an item. Click here to enter text.</p>	<p>N/A</p>	<p>CNS: headache, asthenia, insomnia CV: peripheral edema EENT: pharyngitis, rhinitis, sinusitis GI: abdominal pain, constipation, GU: uti</p>	<ol style="list-style-type: none"> 1. Teach about proper dietary management. 2. Teach pt about weight control, and exercise. 3. Avoid alcohol. 4. Report adverse reactions, muscle pain, malaise and fever.
<p>Atenolol Propranolol</p>	<p>Beta blockers</p>	<p>HTN, Angina pectoris, arrhythmias</p>	<p>PO, IV</p>	<p>Choose an item. Click here to enter text.</p>	<p>Drug compatible with D5W and NSS, and lactated ringer, infusion rate should not exceed 1mg/minute</p>	<p>Cns: fatigue, lethargy, fever CV: hypotension, bradycardia, heart failure, intensification of AV block GI: abdominal cramping, constipation Hematologic: agranulocytosis Respiratory: bronchospasm Skin: rash</p>	<ol style="list-style-type: none"> 1. Continue to take medication, even when feeling better. 2. Take medication with food. 3. Do not suddenly stop medication, it can worsen chest pain, and trigger a heart attack. 4. Advise patients to avoid smoking, alcohol, and illicit drug use.

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

<p>Capoten Captopril</p>	<p>Antihypertensive, Ace inhibitors</p>	<p>HTN, diabetic nephropathy, Heart failure, Left ventricular dysfunction after MI</p>	<p>PO</p>	<p>Choose an item. Click here to enter text.</p>	<p>NA</p>	<p>CNS: dizziness, fainting, headache CV: tachycardia, hypotension, angina GI: abdominal pain, anorexia, constipation Hematologic: leukopenia, agranulocytosis Respiratory: dry, persistent, nonproductive cough</p>	<ol style="list-style-type: none"> 1. Take medication 1 hour before meals, food in gi may reduce absorption. 2. Light-headedness is possible, especially during the first few days of therapy. 3. Rise slowly to minimize this effect and report this effect and to report occurrence to prescriber. 4. Notify prescriber promptly if pregnancy occurs.
<p>Cozaar Losartan</p>	<p>Antihypertensives Angiotensin receptor blockers</p>	<p>HTN, Nephropathy in pt with type 2 diabetes, To reduce risk of stroke in patients with hypertension and left ventricular hypertrophy</p>	<p>PO</p>	<p>Choose an item. Click here to enter text.</p>	<p>NA</p>	<p>CNS: dizziness, asthenia, fatigue CV: edema, chest pain EENT: nasal congestion, sinusitis</p>	<ol style="list-style-type: none"> 1. Avoid salt substitutes, they may contain potassium. 2. Teach women the consequences of becoming pregnant. 3. Do not breast - feed while taking drugs . 4. Report swelling of face, eyes, lips, tongue or any breathing difficulty.

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Imdur Isosorbide dinitrate	Antianginals Nitrates	Acute anginal attacks, Prevention of variceal rebleeding	PO	Choose an item. Click here to enter text.	NA	CNS: headache, dizziness CV: orthostatic hypertension, tachycardia EENT: sublingual	<ol style="list-style-type: none"> 1. Take medication regularly as prescribed. 2. Take a sublingual tablet at the first sign of attack. 3. Place the sublingual up under the tongue until dissolved. 4. Avoid alcohol when taking this medication.
Lasix Furosemide	Antihypertensives Loop diuretics	Acute pulmonary edema edema HTN	PO IV IM	Choose an item. Click here to enter text.	Can dilute with D5W, NSS and Lactated ringer	CNS: vertigo, headache, dizziness, paresthesia CV: orthostatic hypotension, thrombophlebitis with iv administration. EENT: transient deafness, blurred or yellowed vision, tinnitus	<ol style="list-style-type: none"> 1. Take medication in the morning to prevent frequent urination at night. 2. Inform pt for need of potassium or magnesium supplements. 3. Advise pt to report ringing in ears, severe abdominal or sore throat and fever immediately. 4. Stand slowly to prevent dizziness and to limit alcohol.
Digoxin	Cardiac Glycosides	Heart failure Atrial Fibrillation	PO IV	Choose an item. Click here to enter text.	Can dilute with d5w, nss, or sterile water for injection, infuse drug slowly over at least 5 minutes	CNS: agitation, fatigue, generalized muscle weakness CV: arrhythmias, heart block	<ol style="list-style-type: none"> 1. Teach pt how to take v/s, before taking medication. 2. Teach pt to report pulse rate less than 60 beats/min . 3. Report adverse reactions, visual disturbances may be indicators of toxicity. 4. Encourage pt to eat a consistent amount of potassium rich foods.

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

						EENT: blurred vision, diplopia, light flashes GI: anorexia, n/v/d	
Cardizem Diltiazem	calcium channel blockers	To manage Prinzmetal or variant angina or chronic stable angina pectoris, HTN, Atrial fibrillation	PO IV	Choose an item. Click here to enter text.	Can dilute with d5w, nss, and d5w mixed with nss	CNS: headache, dizziness CV: edema, arrhythmia, AV block, bradycardia, heart failure GI: nausea, constipation, abdominal discomfort Hepatic: acute hepatic injury	<ol style="list-style-type: none"> 1. Take medication as prescribed. 2. Avoid hazardous activities during the start of therapy. 3. Take medication as prescribed. 4. Report any signs of adverse reaction.
Coumadin Warfarin	Anticoagulant	Pulmonary embolism, DVT, MI, AFIB,	PO IV	Choose an item. Click here to enter text.	Draw up with sterile water, 2.7 ml	CV: vasculitis GI: abdominal pain, diarrhea, flatulence Hematologic: hemorrhage Hepatic: hepatitis	<ol style="list-style-type: none"> 1. Do not take with cranberry juice, may increase the risk of bleeding. 2. Have blood drawn and levels checked regularly before taking medication. 3. Avoid OTC products containing ASA, and other salicylates. 4. Watch for signs of abnormal bleeding, bruising and report immediately to the prescriber.

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Plavix Clopidogrel	Antiplatelet s	To reduce thrombotic events, st segment elevation acute MI	PO	Choose an item. Click here to enter text.	N/A	<p>CNS: confusion , fatal intracranial bleeding</p> <p>CV: hypotension</p> <p>EENT: epistaxis, rhinitis, taste disorder</p> <p>GU: hemorrhage, abdominal pain</p> <p>Hematologic: thrombotic thrombocytopenic purpura</p> <p>Resp; bronchospasm</p>	<ol style="list-style-type: none"> 1. May take longer than usual to stop bleeding. 2. Report to prescriber unusual bleed or bruising occurs. 3. Inform all health care and dentist about medication before any procedures. 4. May take with or without food.
-----------------------	-------------------	--	----	--	-----	---	--