

| | |
|--|--|
| <p>Patient #1 Diagnosis/Chief Complaint</p> <p>Tibia Fracture</p> | <p>Patient #1 Teaching</p> <ol style="list-style-type: none"> 1. Teach patient how to use crutches 2. Teach patient to elevate injured leg to reduce swelling 3. Teach patient how to clean injured leg without disturbing the cast <p>Discharge planning/Community Resources</p> <ol style="list-style-type: none"> 1. Physical therapy 2. Follow-up appointment to remove cast 3. Home health to help bathe patient if needed |
| <p>Patient #2 Diagnosis/Chief Complaint</p> <p>Appendicitis</p> | <p>Patient #2 Teaching</p> <ol style="list-style-type: none"> 1. Teach parent the minimum and maximum pain medication doses 2. Teach patient to limit sports or strenuous activities for 2 weeks 3. Teach parent to call PCP immediately at sight of swelling, oozing, or increased pain <p>Discharge Planning/Community Resources</p> <ol style="list-style-type: none"> 1. Consult home health for surgical site cleaning at home if necessary 2. Schedule follow-up appointment with PCP 3. Consult dietician to consult with patient and parent about post-op diet |
| <p>Patient #3 Diagnosis/Chief Complaint</p> <p>Post-Ictal Seizure</p> | <p>Patient #3 Teaching</p> <ol style="list-style-type: none"> 1. Teach patient and parent to take seizure medication daily 2. Teach parent how to keep patient safe during seizure episode 3. Teach patient to not swim alone |

| | |
|--|---|
| | <p>Discharge Planning/Community Resources</p> <ol style="list-style-type: none">1. Get patient involved in a seizure support group2. Send patient and parent home with seizure log to keep track of episodes3. Send patient home with a medical alert bracelet |
|--|---|