

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
Potassium Chloride 20 mEq in 1/2 NS + KCL 20 mEq/L 1,000 mL, 100 mL/hr	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input checked="" type="checkbox"/>	It is an electrolyte replenisher. Used to treat or prevent low blood levels of potassium.	Electrolyte panel	Can cause diarrhea, flatulence, nausea, vomiting, cardiac arrest, hyperkalemia, GI ulcer.

Student Name: Kambree Irvin		Unit: 3 N	Patient Initials: <a href="#">Click here to enter text.</a>		Date: 1/4/2022	Allergies: NKA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Nafcillin Sodium	Antibiotic	Used to treat many different types of infections.	2g in NS 50 mL, IVPB, Q4h		- concentration of 40mg, 100 ml/hr	Injection site extravasation, hypokalemia, cdiff, bone marrow depression, agranulocytosis, tubulointersitial nephritis	1. Instruct patient to report severe diarrhea 2. Monitor IV site for redness, swelling, pain 3. Monitor CBC, liver function, & renal function 4. If self administered, teach proper injection technique, placement of injection, and rotation of injection sites.
Trazodone	Antidepressant	Used to treat major depressive disorder.	50 mg/tablet, PO, QHS	This medication isn't recommended for use for adolescents.		Constipation, diarrhea, nausea, vomiting, confusion, fatigue, cardiac dysrhythmia, seizure,	1. Possibly call HCP, as trazodone is not recommended in children or adolescents. 2. Instruct patient/family to report unusual changes in behavior or suicidal thoughts 3. Avoid mental alertness activities as drug can cause dizziness, confusion, somnolence

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Kambree Irvin		Unit: 3 N	Patient Initials: Click here to enter text.		Date: 1/4/2022	Allergies: NKA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
						serotonin syndrome, suicide, priapism	4. Take after a meal or snack
Sertraline	Antidepressant	Used to treat major depressive disorder, OCD, panic disorder, social anxiety, or PTSD.	100 mg/tablet, PO, QHS	Click here to enter text.	Click here to enter text.	Constipation, indigestion, dizziness, insomnia, fatigue, hyponatremia, hemorrhage, seizure, depression, mania, serotonin syndrome	<ol style="list-style-type: none"> <li>1. Monitor for mental status changes, neuromuscular changes, autonomic instability</li> <li>2. Avoid activities requiring mental alertness until effects are realized</li> <li>3. Inform patient/family against sudden discontinuation</li> <li>4. Educate patient of the side effects including sweating, tremor, nausea, dry mouth, and dizziness</li> </ol>
Fluticasone Propionate	Adrenal Glucocorticoid	Used to treat nasal congestion, sneezing, runny nose, and itchy watery eyes.	1 spray, 50 mcg, both nares daily		Click here to enter text.	Osteoporosis, headache, bleeding from nose, cough, nasopharyngitis, sinusitis, throat irritation, upper resp infection,	<ol style="list-style-type: none"> <li>1. Rinse mouth with water after to reduce risk of oral candidiasis.</li> <li>2. Monitor for symptoms that do not improve or worsen.</li> <li>3. Administer at same time every day and do not use more than twice every 24 hours.</li> </ol>

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

<b>Student Name:</b> Kambree Irvin		<b>Unit:</b> 3 N	<b>Patient Initials:</b> <a href="#">Click here to enter text.</a>		<b>Date:</b> 1/4/2022	<b>Allergies:</b> NKA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
						oral candidiasis, pneumonia	4. Medication can lower blood cells that help fight infections, avoid being around sick people or chicken pox/measles.
Clonidine	Antihypertensive	Lowers blood pressure by decreasing certain chemicals in the blood. Used to treat hypertension. The Kapvay brand is used to treat ADHD.	0.4mg/2 tablets, PO, Q24hr	<input type="text" value=""/>	<input type="text" value=""/> here to enter text.	Contact dermatitis, erythema, pruritus, xerostomia, dizziness, headache, somnolence, fatigue	<ol style="list-style-type: none"> <li>1. Possible withdrawal symptoms if sudden discontinuation occurs</li> <li>2. If vomiting occurs, notify HCP. This is especially important for children.</li> <li>3. This medication may cause brain fog.</li> <li>4. It is best to avoid over the counter medications that contain alcohol while taking this medication.</li> </ol>

## Pediatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: Kambree Irvin		Unit: 3 N	Patient Initials: Click here to enter text.		Date: 1/4/2022	Allergies: NKA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP – List diluent solution, volume, and rate of administration  IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Methylphenidate	CNS Stimulant	Affects chemicals in the brain and nerves that contribute to hyperactivity and impulse control. Used to treat ADD & ADHD.	5mg/ tablet, PO Q24hr	Click here to enter text.	Click here to enter text.	Erythema, decrease in appetite, nausea, vomiting, headache, nasal congestion, mania, raynauds disease, drug dependence	<ol style="list-style-type: none"> <li>Misuse may cause addiction, overdose, or death. Inform family to store where others cannot get to it.</li> <li>May be taken with or without food, but take it the same way each time</li> <li>Inform doctor of other medications, especially antidepressants. As they can interact with one another.</li> <li>Monitor for aggressive behavior and hostility.</li> </ol>
Furosemide	Loop Diuretic	Used to treat fluid retention and high blood pressure.	40 mg/4mL, IVP, once	Click here to enter text.	Click here to enter text. um Chloride or undiluted, 1-2 minutes	Hyperuricemia, Hypomagnesemia, spasm of bladder, orthostatic hypotension, pancreatitis, thrombocytopenia	<ol style="list-style-type: none"> <li>Avoid giving close to bedtime as it can cause frequent urination</li> <li>Monitor for bruising or bleeding</li> <li>Have child avoid actions or tasks that call for alertness</li> <li>Monitor for confusion</li> </ol>
Albumin	Volume Expander	Replace blood volume loss resulting from trauma.	30mg/120mL, IVPB, once	Click here to enter text.	Click here to enter text. /hr, 0.25 mg	Hypovolemia, Hypersensitivity reaction	<ol style="list-style-type: none"> <li>Monitor vitals, electrolytes, and kidney function</li> <li>Drink plenty of fluids while treated with albumin</li> </ol>

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Kambree Irvin		Unit: 3 N	Patient Initials: <a href="#">Click here to enter text.</a>	Date: 1/4/2022	Allergies: NKA		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							3. Labs will need to be checked often 4. Monitor for shortness of breath, swelling in arms or legs, and weight gain
							1. <a href="#">Click here to enter text.</a> 2. <a href="#">Click here to enter text.</a> 3. <a href="#">Click here to enter text.</a> 4. <a href="#">Click here to enter text.</a>
Patient 2 - Right INT, NKA	<a href="#">Click here to enter text.</a>	<a href="#">Click here to enter text.</a>	<a href="#">Click here to enter text.</a>	<a href="#">Click here to enter text.</a>	<a href="#">Click here to enter text.</a>	<a href="#">Click here to enter text.</a>	1. <a href="#">Click here to enter text.</a> 2. <a href="#">Click here to enter text.</a> 3. <a href="#">Click here to enter text.</a> 4. <a href="#">Click here to enter text.</a>
Amoxicillin susp.	Antibiotic	Penicillin that fights bacteria. Used for many different types of infections.	512mg/6.4mL, PO, BID	<a href="#">Click here to enter text.</a>	<a href="#">Click here to enter text.</a>	Rash, diarrhea, vomiting, nausea, headache, Stevens Johnsons syndrome, cdiff	1. Shake oral suspension before measuring a dose 2. Administer at the same time each day 3. Do not skip a dose. Skipping can increase risk of infection that is resistant to medication. 4. Monitor for diarrhea, could be a sign of a new infection
Ethosuximide	Anticonvulsant	Used to treat absence	375mg/7.5ml, PO	<a href="#">Click here to enter text.</a>	<a href="#">Click here to enter text.</a>	Loss of appetite, nausea, ataxia, headache,	1. Monitor skin reactions such as a spreading red rash, blistering

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

<b>Student Name:</b> Kambree Irvin		<b>Unit:</b> 3 N	<b>Patient Initials:</b> Click here to enter text.		<b>Date:</b> 1/4/2022	<b>Allergies:</b> NKA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
		seizures.	QHS	enter text.		aplastic anemia, pancytopenia, leukopenia, lupus, seizure, depression	<ol style="list-style-type: none"> <li>2. Report s/s of blood dyscrasias (sore throat, fever, bruising, malaise)</li> <li>3. Avoid activities requiring mental alertness.</li> <li>4. Report unusual changes in behavior, depression, suicidal ideation.</li> </ol>
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	<input style="width: 100px; height: 20px;" type="text"/> here to enter text. Click here to enter text.	Click here to enter text.	Click here to enter text.	<ol style="list-style-type: none"> <li>1. Click here to enter text.</li> <li>2. Click here to enter text.</li> <li>3. Click here to enter text.</li> <li>4. Click here to enter text.</li> </ol>
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	<input style="width: 100px; height: 20px;" type="text"/> here to enter text. Click here to enter text.	Click here to enter text.	Click here to enter text.	<ol style="list-style-type: none"> <li>1. Click here to enter text.</li> <li>2. Click here to enter text.</li> <li>3. Click here to enter text.</li> <li>4. Click here to enter text.</li> </ol>