

1. _____ - smell
2. _____ - Visual acuity
3. _____ - Opening of eyelids, eye movement upward/ medial, upward/lateral,
medial, downward/ lateral)
4. _____ - Eye movement (downward/ Medial)
5. _____ - Facial sensation, chewing movement
6. _____ - Eye movement (lateral)
7. _____ - Facial muscle movement (except chewing muscles) and eyelid closing
8. _____ - Hearing and balance
9. _____ - Taste on the posterior third of the tongue
10. _____ (Palate muscles) and swallowing
11. _____ - Shoulder shrug
12. _____ - Tongue movement
13. I can lead to Ineffective communication

14. I can lead to decrease interaction, withdrawal, suspicion, loss of self-esteem and insecurity
15. I affect the outer and middle ear
16. I affect the inner ear and affect nerve pathways
17. I affect the inner, middle, and outer ear
18. I am an assessment tool used to look at the ear
19. I am an assessment test used that compares air conduction with bone conduction
20. I am normal when air conduction is louder than bone conduction
21. I am abnormal when bone conduction is louder than air conduction
22. I am an assessment test when conducted I will strike a 512hz tuning fork and place on my forehead
23. I am normal when tuning fork is on my forehead and can equally be heard in both ears
24. I am abnormal when a tuning fork is on my forehead and the sound is different in my two ears.
25. I am the medical term of ringing of the ears
26. I am the medical term for the room is spinning

27. I am a symptom that occurs when there is fluid in the ear compressing the 8th cranial nerve
28. I am the reason the patient is a high fall risk
29. I can cause a patient grief and depression and may also lead to suicidal ideation
30. I am caused by Otosclerosis
31. I am caused by otitis media with effusion
32. I am caused by Poor Eustachian tube function
33. I am caused by impacted cerumen (earwax)
34. I am caused by tumors in the ear
35. I am caused by objects in the ear (such as a Lego)
36. I hear better in noisy environment (bone better than air conduction)
37. My treatment plan is to either treat the cause or wear a hearing aid.
38. I am the most common type of permanent hearing loss
39. I cannot be fixed by surgery or medicine
40. I am caused by illness or genetics

41. I am caused by loud noises like concerts, gun firing, construction, factory work etc.

42. I am caused by ototoxicity

43. I am Vancomycin I cause what kind of damage

44. I am Gentamycin I cause what kind of damage

45. I am Cisplatin I cause what kind of damage

46. I am Aspirin I cause what kind of damage

47. I am Furosemide I cause what kind of damage

48. I am Quinine I cause what kind of damage

49. I am caused by presbycusis (aging)

50. I am caused by dilation of the endolymphatic system by overproduction or decrease

reabsorption of endolymphatic fluid.

51. My signs and symptoms include: Vertigo, Tinnitus, and Fluctuating hearing loss

52. My interventions include: prevent injury during Vertigo attacks, provide bed rest in a quiet

environment, provide assistance with walking, instruct the patient to move their head slowly to prevent

worsening of the Vertigo, initiate sodium and fluid restrictions as prescribed, instruct the patient to stop smoking, instructed patient to avoid watching television because the flickering of lights may exacerbate symptoms, allow the patient to rest; And control Vertigo nausea in vomiting, mild diuretics may be prescribed to decrease endolymphatic volume, inform the patient about vestibular rehabilitation as prescribed.

53. My surgical intervention includes a resection of the vestibular nerve or a total removal
54. My post-operative interventions- packing and dressings on the ear. Speak to the patient on the side of the unaffected ear. Perform neurological assessments. Maintain safety. Assess when ambulating. Encourage the patient to use a bedside commode rather than ambulating to the bathroom. Administer anti- vertigo and anti-emetic medications as prescribed.
55. For me you should assess for early detection to prevent permanent damage
56. For me you should educate patients and have them teach back to ensure understanding
57. I must be taught about adequate nutrition and hydration to ensure that I can maintain oil and sebaceous glands to maintain hearing.
58. I must be taught about medication regimens

- 59. I must be taught about surgery if necessary
- 60. You should never shout at me
- 61. You should speak low and slow
- 62. You can write out what you are trying to say
- 63. I would like some help with the cost of my hearing aids