

Maternal Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: NKDA

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Oxytocic Agent	Increases the amplitude and frequency of uterine contractions	0.5 to 1 mU/ min, IV, Q30-60	Y N	IVPB- 0.5 to 1 mU/min and gradually increase 1 to 2mU/min every 30 to 60 minutes.	Nausea, vomiting, cardiac dysrhythmia, fetal bradycardia, water intoxication	1. BBW- Not indicated for elective induction of labor 2. Patient should not drink any fluids because of risk of water intoxication. 3. Teach patient how the oxytocin will affect their contractions 4. Report any signs of headache, dizziness, and intense pain.
Magnesium Sulfate	Sulfate	Prevents seizures	IV- 4 to 5g over 3 to 4 minutes IM- 10g , 5 in each buttock	Y N	IV- mix in 250 mL of D5W or NS	Flushing, sweating, heart block, hypotension, myocardial dysfunction, shock, hypocalcemia, hypothermia	1. Teach patient signs and symptoms of magnesium toxicity. 2. Precaution in patients with renal insufficiency because can cause magnesium intoxication. 3. Precaution in giving medication for more than 5 to 7 days because can lead to hypocalcemia and bone abnormalities in the fetus. 4. Contraindicated in mothers with toxemia of pregnancy during the 2 hours preceding delivery.

Maternal Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: NKDA

Meperidine	Narcotic analgesic	Pain relief	0.2 to 0.5 mg/kg IV 50 to 150 mg orally/subQ/IM Q3-4	Y N	N/A	Sweating, nausea, vomiting, dizziness, lightheadedness, cardiac arrest, hypotension	<ol style="list-style-type: none"> 1. BBW- can lead to addiction, abuse, and misuse. 2. Avoid any activities that require any mental alertness or coordination. 3. Report severe constipation. 4. Avoid alcohol and other CNS depressants.
Promethazine	Antihistamine	Treats allergies, nausea and vomiting	25 to 50 mg PO/rectally 25 to 50 mg deep IM 25 mg/mL at 25mg/min	Y N	IV- 25 mg/ml at 25 mg/min	Dermatitis, phototoxicity, xerostomia, CNS depression, prolonged QT interval	<ol style="list-style-type: none"> 1. BBW- can cause severe chemical irritation and damage to tissues 2. Avoid any prolonged exposure to the sun. Wear hats and sunscreen when out in the sun. 3. Avoid drinking while taking this medication. 4. Report any signs of sweating, fever, stupor, unstable blood pressure, muscular rigidity, and autonomic dysfunction (EPS/NMS)
Calcium Gluconate	Antidotes	Treats conditions arising from calcium deficiencies	1000 to 2000 mg, IV, every Q6	Y N	N/A	Abnormal taste in mouth, constipation, flatulence, swollen abdomen, cardiac arrest, bradyarrhythmia	<ol style="list-style-type: none"> 1. May cause tingling sensation, chalky taste, or heat waves. 2. Take with a whole glass of water. 3. Should take oral powder with food. 4. Contraindicated in patients with ventricular fibrillation.

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
--	-----------------	-------------------	-------------------------------------	---------------------------------

Maternal Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: NKDA

	Isotonic/ Hypotonic/ Hypertonic			
--	---------------------------------	--	--	--

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	Vitamins	Used to treat vitamin K deficiency and to treat certain bleeding or blood clotting problems.	2.5 to 25 mg orally/ IM/iv? subQ	Y N	N/A	Cardiac arrest, shock, cutaneous hypersensitivity, metabolic acidosis, gasping for breath, CNS depression	<ol style="list-style-type: none"> 1. BBW- can cause fatalities with severe reactions. 2. Report any rashes. 3. Contraindication for any hypersensitivity to any component of this medication. 4. Precaution in patients with preexisting liver disease.
Erythromycin Ophthalmic Ointment	Antibiotic	Treat bacterial infection of the eyes.	1 cm ribbon, up to 6 times a day	Y N	N/A	Diarrhea, stomach cramps, loss of appetite, prolonged QT interval, Stevens-Johnson syndrome	<ol style="list-style-type: none"> 1. Do not use with astemizole, cisapride, pimozone, terfenadine, or lovastatin. 2. If having severe diarrhea report to your PCP before taking any antidiarrhea medication. 3. Contraindicated in patients with hypersensitivity of erythromycin. 4. Precaution in patients with hepatic impairment because of increased risk of toxicity.

Maternal Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: NKDA

Engerix B	Recombina nt	Prevents hepatitis B in adults	10 mcg/ 0.5 mL IM Q 6 months	Y N	N/A	Injection site disorder, nausea, diarrhea, astenia, fatigue, anaphylaxis	1. Contraindicated in patietns with previous reaction to any hepatitis B containing vaccine or yeast. 2. Precation in patients who are sensitive to latex. 3. Precaution in patients who are ill. 4. Administer within 12 hours of birth.
Hepatitis B Immune Globulin	Immune Serum	Prevents Hepatitis B	0.06 mL/kg, IM,	Y N	N/A	Erythema at injection site, injection site pain, headache, malasie, ecchymosis, decreased WBCs	1. Avoid any live vaccines for 3 months after receiving this drugs. 2. Can interfere with with blood glucose monitoring systems. 3. Human plasma products carry a risk of transmitting infectious agents. 4. Precaution in advanced age because of thrombotic events.
				Y N			1. 2. 3. 4.