

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**Allergies:** _____

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Oxytocic hormones	Induce labor or strengthen contractions		Y N		Irregular heart rate, severe headache, confusion, N/V Excessive bleeding after birth	1. Do not give if pelvis is not sufficient for labor 2. Do not give if active herpes virus 3. Do not give if mom has high BP 4. Monitor contractions and fetal activity
Magnesium Sulfate		Prevent preterm labor or prevent seizures		Y N		N/V, thirst confusion, sweating	1. used longer than 5-7 days can cause low calcium levels in baby 2. caution magnesium toxicity 3. monitor vital signs 4. monitor for signs dehydration
Meperidine	Opioid	Treats moderate to severe pain		Y N		Slow heart rate, shallow breathing, severe drowsiness, confusion, constipation	1. misuse can cause addiction overdose or death 2. do not use if MAOI has been used in past 14 days 3. do not take if you have asthma 4. can pass into breast milk
Promethazine	phenothiazines	Treats nausea and pain after surgery		Y N		Drowsiness, confusion, irregular heartbeats,	1. can impair thinking or reactions 2. avoid getting up too fast 3. limit sun exposure

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

						weakness	4. best taken at night
Calcium Gluconate		Treats or prevents low calcium levels		Y N		Weakness, headache, bone pain, dizziness, slow heartbeat	1. tell doctor of all medications and vitamins 2. can affect certain labs 3. can cause risks to baby if breastfeeding 4. can cause tissue damage if leaked from vein

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione		Treats vitamin k deficiency		Y N		Dizziness, flushing, sweating, pain or swelling at injection site	5. Can cause serious side effects or death to premature babies 2. tell your dentist before appointment 3. protect medication from light 4. call your doctor if you have an unusual or unpleasant taste in your mouth
Erythromycin Ophthalmic Ointment	Antibiotic	Fights bacteria		Y N		Eye redness, mild irritation	1. keep eye closed 1-2 minutes after administration 2. avoid driving after 3. do not touch tip of tube or place on eye 4. skipping or stopping doses can increase risk of infection
Engerix B		Prevent hep B		Y N		Headache, tiredness, redness/pain at site	1. cannot treat an active infection 2. recommended for adults at risk for hep B 3. do not take if allergic to yeast 4. series of 2-4 shots

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

Hepatitis B Immune Globulin	Human plasma	Prevent hep B in pts receiving liver transplant or who have been exposed		Y N		Fever, mouth sores, lightheadedness liver problems, chest pain, symptoms of blood clot	<ol style="list-style-type: none"> 1. this does not provide long term protection, do not take with live vaccine 2. can increase risks of blood clots 3. not known if this can harm a baby or passes in breast milk 4. given to baby within 12 hours of birth if mother was infected with hep B
				Y N			<ol style="list-style-type: none"> 1. 2. 3. 4.