Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
Click here to enter text.	Isotonic ☐ Hypotonic ☐ Hypertonic ☐	Click here to enter text.	Click here to enter text.	Click here to enter text.

Student Name:	Unit: PMH SM	Patient Initials:	Date: 4/5/22	Allergies:
Click here to enter text. Keslee Jenkins	Click here to enter text.	Click here to enter text.	Click here to enter a date.	Click here to enter text.

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Hydroxyzine pamoate	Antihistamine	Treats anxiety and itching caused by allergies	25mg PO PRN q6 for anxiety	Yes	N/A	Dry mouth, somnolence, prolonged QT interval, dizziness, blurred vision, constipation	1. Avoid any alcohol and other CNS depressants. 2. Do not operate heavy machinery until you know how the medication affects you. 3. Use birth control because could be harmful to a baby. 4. Take with food to avoid an upset stomach.
lorazepam	Benzodiazapine	Treats anxiety, insomnia, sleep difficulty due to anxiety or stress, and for alcohol withdrawal	1-2 mg PO PRN q4 hours for alcohol withdrawal	Yes	IVPB- dilute with equal volume of compatible solution immediately prior to use, mix gently. Run at a rate of 2mg/minute	Asthenia, dizziness, sedated, unsteadiness, depression, acidosis, delirium	1. BBW- can lead to abuse, misuse and addiction of benzos. 2. Avoid pregnancy as the risk to the fetus are unknown. 3. Do not quit taking medication suddenly. 4. Teach patient about the misuse of benzos and the symptoms and risks.
Ziprasidone	psychotropic agent	↑ dopamine + serotonin in the brain	40mg PO BID	Yes	N/A	rash, weight gain, weight loss, dry mouth, anxiety, dizziness, increased muscle tone, tremor, abnormal vision	1. BBW - elderly pts with dementia related psychosis are at ↑ risk for death 2. Take with food 3. Avoid activities that require mental alertness until drug is realised 4. Teach sxs of neuroleptic maglinant syndrome or tardive dyskinesia.

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text.		**Unit:** Click here to enter text.	**Patient Initials:** Click here to enter text.		**Date:** Click here to enter a date.	**Allergies:** Click here to enter text.	
Generic Name	**Pharmacologic Classification**	**Therapeutic Reason**	**Dose, Route & Schedule**	**Correct Dose? If not, what is correct dose?**	**IVP – List diluent solution, volume, and rate of administration** **IVPB – List concentration and rate of administration**	**Adverse Effects**	**Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)**
acetaminophen	Analgesic/ antipyretic	Reduces pain	500 mg PO q6 for pain of 1-4 on pain scale	Yes	N/A	Pruritus, constipation, nausea, insomnia, agitation	1. It is not safe to take more than 4000 mg in a 24 hour period. 2. Take with a full glass of water. 3. Avoid taking with alcohol. 4. Avoid pregnancy as the risk to the fetus are unknown.
olanzapine	Atypical antipsychotic	Rebalances dopamine and serotonin in the brain.	5 mg PO q6 for agitation	Yes	N/A	Peripheral edema, hyperglycemia, increased appetite, tremor, increased weight, dizziness	1. BBW- risk of death is increased in elderly patients with dementia related psychosis. 2. Report any rashes, fever, or other symptoms of drug reactions. 3. Avoid alcohol. 4. Advise patient to symptoms of orthostatic hypotension.
ibuprofen	NSAIDS	Pain reliver	400 mg PO q6 for pain of 1-4 on pain scale	Yes	N/A	Hypotension, hypernatremia, rash, flatulence, thrombocytosis, urinary retention, cardiac arrest, Heart failure	1. BBW- can cause increased risk for cardiovascular events like stroke. 2. Avoid taking any extra NSAIDS or aspirin at the same time. 3. Take with food or milk to avoid GI upset. 4. Report any symptoms of skin reaction.

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text. Keslee Jenkins	Unit: SIM Click here to enter text.	Patient Initials: Click here to enter text.	Date: 4/5/22 Click here to enter a date.	Allergies: Click here to enter text.

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
haldol	antipsychotic	Helps control psychotic symptoms	10 mg IM PRN q8 for psychotic symptoms /agitation	Yes	N/A	Hypotension, constipation, dry mouth, akathisia, blurred vision, hyponatremia, pneumonia, tardive dyskinesia	**1.** BBW- avoid use in elderly patients iwth dementia related psychosis. **2.** Avoid sudden discontinuation of this drug. **3.** This drug may impair heat regulation which can lead to increased core temperature. **4.** Report any jerky muscle movements, tongue thrusting, ticks, or random movements of your extremities.
diphenhydramine	antihistamine	Treats sides effects of certain psychiatric drugs such as involuntary movements and muscle stiffness	50 mg IM q8 PRN for psychotic symptoms /agitation	Yes	N/A	Dry mouth, dizziness, dyskinesia, sedation, somnolence, thick sputum	**1.** Avoid activities that require mental alertness or coordination until drug effects are realized. **2.** Avoid alcohol. **3.** Avoid concomitant use of MAO inhibitors or CNS depressants. **4.** This medication may cause blurred vision and may impair your thinking or reactions.
Prazosin quinazoline derivative	treats PTSD	1 mg PO QHS	yes	N/A	Orthostatic hypotension, palpations, nausea, lethargy, somnolence, headache, dizziness	1. Avoid alcohol 2. May experience syncope or loss of consciousness 3. Do not stop taking suddenly 4. Rise slowly from a sitting/laying position because can cause orthostatic hypotension.	

Adopted: August 2016

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text.		**Unit:** Click here to enter text.	**Patient Initials:** Click here to enter text.		**Date:** Click here to enter a date.	**Allergies:** Click here to enter text.	
Generic Name	**Pharmacologic Classification**	**Therapeutic Reason**	**Dose, Route & Schedule**	**Correct Dose? If not, what is correct dose?**	**IVP – List diluent solution, volume, and rate of administration** **IVPB – List concentration and rate of administration**	**Adverse Effects**	**Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)**
risperdal	antipsychotic	Rebalances dopamine and serotonin in the brain.	1 mg tablet PO BID	Yes	N/A	Rash, increased weight, excessive salivation, upper abdominal pain, dry mouth, dizziness, tremor, sedation, increased appetite	**1.** BBW- risk of death is increased in elderly patients with dementia related psychosis. **2.** Avoid activities that require mental alertness or coordination until drug effects are realized. **3.** Rise from a sitting/lying position slowly because can cause orthostatic hypotension. **4.** This drug may impair heat regulation which can lead to increased core temperature.
nicotine	Stimulant of autonomic ganglia	Treats smoking and withdrawal syndrome.	21 mg transdermal patch daily	Yes	N/A	Skin irriration, nicotine withdrawal, dizziness, headache, insomnia, cardiac dysrhythmia	**1.** Transdermal patch may cause application site reactions. **2.** Report any signs and symptoms of cardiovascular events. **3.** Do not use other nicotine containing products while using the patch. **4.** May cause tobacco withdrawal symptoms such as insomnia, anxiety, and restlessness.