

Student Name: Kambree Irvin_____

Unit:___SIM___

Pt. Initials: _____

Date: ___5/9/22_____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Pituitary Hormone, Uterine Stimulant	Used to induce labor or strengthen contractions or control bleeding after childbirth.	IV, determined by uterine response	Y N	Dilute 10 unites in 1000 mL of NS or LR	Nausea, vomiting, cardiac dysrhythmia, increased blood pressure, neonatal jaundice, hemorrhage, hematoma, rupture of uterus	1. Black box: since available data are inadequate to evaluate risks to benefits considerations, oxytocin is not indicated for elective induction of labor. 2. Follow caregiver's instructions about drinking or restricting fluids 3. Monitor for fast, slow, or uneven heart rate 4. Watch closely for excessive bleeding long after childbirth
Magnesium Sulfate	Anti-inflammatory, laxative, musculoskeletal agent	Important to muscles and nerves, increases water in the intestines, used to relieve constipation, can be helpful to relieve minor sprains	IM or IV, PO	Y N	IV- dilute with D5W or NS to a concentration of 20% or less, max rate is 150mg/min	Flushing, sweating, heart block, hypotension, hypocalcemia, hypothermia, CNS depression	1. Drink plenty of liquids while taking 2. Monitor for a sudden change in bowel habits that has lasted 2 weeks or longer 3. Tell doctor if breastfeeding, as it is unsure if this drug could harm a nursing baby 4. Stop medication if rectal bleeding occurs

Student Name: Kambree Irvin_____

Unit: SIM_____

Pt. Initials: _____

Date: 5/9/22_____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

Calcium Gluconate	Parental mineral-trace mineral	Used to treat or prevent low calcium levels	IV. PO	Y N	Dilute in D5W or NS	Constipation, swollen abdomen, cardiac arrest, hypotension, vasodilation, prostate cancer	1.Take oral powder with food or liquid 2.Monitor for drug-drug interactions as there are multiple 3.Monitor lab tests closely 4.Use in care with newborns, tell doctor if you are pregnant or plan to get pregnant, or if you are breast-feeding
-------------------	--------------------------------	---	--------	--------	---------------------	---	---

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	Vitamin K	Treats certain bleeding or blood clotting problems	IV, PO, Subq	Y N	IV- dilute in NS or D5W in a minimum of 50 mL over 20 minutes or a rate to not exceed 1mg/min	Cardiac arrest, shock, metabolic acidosis, CNS depression, gasping for breath, respiratory arrest	1.Monitor for rashes 2. Injection may contain ingredient that can cause serious side effects or death in very young and premature babies. 3.May need to receive frequent blood tests 4.Black box: severe reactions, including fatalities, have occurred during or after injection, monitor patient extremely closely
Erythromycin Ophthalmic Ointment	Antibiotic	Fights bacteria and treats bacterial infections of the eyes	Topical	Y N		Nausea, vomiting, abdominal pain, diarrhea, anorexia,	1.Do not use is a viral or fungal infection of the eye is present 2.Wash hands before applying ointment 3.Tilt head back slightly and pull-down lower eyelid to create a small pocket, avoid touching

