

Student Name:  Mary Lasap Unit:  Sim OB  Pt. Initials: \_\_\_\_\_Date:  06/01/2022 **Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Oxytocics	Induced labor and control postpartum bleeding		Y N		increased uterine motility, B/p changes, uterine rupture, water intoxication	1. Inform pt. to expect contractions like period cramp after administration. 2. Monitor fetal maturity, presentation, pelvic adequacy prior to administration. 3. Monitor V/S, I & O, lung sounds 4. Stop infusion and turn pt. left side if significant fetal heart rate develops.
Magnesium Sulfate	Minerals/ electrolytes	Resolution of eclampsia		Y N	IVPB: 4-6 g over 15-20 min	Magnesium toxicity, flushing diarrhea, hypotension, respiratory depression	1. Report if Magnesium toxicity occurs. 2. Monitor BP, Urine output, respiratory rate, and test deep tendon reflexes. 3. Stop the infusion if toxicity occurs and notify provider. 4. Have resuscitation equipment and Calcium gluconate at bedside.
Meperidine	Opioid agonists	Opioid analgesics		Y N		Confusion, HA, sedation, N/V, hypotension, respiratory depression, constipation	1. Change position slowly to minimize OH. 2. Avoid alcohol or CNS depressants. 3. Monitor Bp, pulse, and respirations before and after throughout the therapy. 4. Take med as directed, do not increase dose without consulting providers.
Promethazine	Phenothiazines	Antiemetics, antihistamines		Y N		Confusion, disorientation, sedation, dry mouth, fatigue,	1. May cause drowsiness, avoid driving or other activities that require alertness. 2. Frequent mouth rinses, sugarless candy or gum may decrease dry mouth.

Student Name: \_\_\_\_\_

Unit: \_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: \_\_\_\_\_

**Newborn Medication Worksheet – Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

						constipation, photosensitivity	3. Use sunscreen and protective clothing to prevent photosensitivity reactions. 4. Avoid alcohol or CNS depressants.
Calcium Gluconate	Antacid	Antidote for magnesium toxicity, mineral and electrolyte replacement		Y N		Phlebitis, hypercalcemia, arrhythmias, constipation, nausea, constipation	2. Take med as directed, take missed doses as soon as possible unless time for next dose. 2. Risk for constipation, increase fluid intake and mobility to prevent constipation. 3. Avoid excessive use of tobacco and beverage containing alcohol or caffeine. 4. Avoid taking calcium-containing supplements during therapy.

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	Fat soluble vitamins	Antidotes, vitamins		Y N		GI upset, rash, flushing, allergic reactions, unusual taste	3. Take med as directed, take missed doses as soon as possible unless time for next dose. 2. Inform pt. to notify if unusual bleeding, nosebleed, excessive menstrual blood flow occurs. 3. Use a soft toothbrush, do not floss, shave with electric razor. 4. Monitor pulse and BP frequently, side effects and adverse reactions.
Erythromycin Ophthalmic Ointment	Macrolides	Antibiotics, treat infections		Y N		Rash, itching, burning,	1. Notify physician if signs of superinfection reactions occur.

Student Name:  Mary Lasap

Unit:  Sim OB  Pt. Initials: \_\_\_\_\_

Date:  06/01/2022

**Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

		of the eyes				swelling, changes in vision, redness	<ol style="list-style-type: none"> <li>2. May blur the visions for a few minutes.</li> <li>3. Use a directed, wash hands before applying the medication.</li> <li>4. Avoid touching applicator tip to any surface.</li> </ol>
Engerix B	Viral	Vaccines for Hepatitis B virus		Y N		Pain/swelling/ redness at the injection site, fever, HA, N/D, loss of appetite	<ol style="list-style-type: none"> <li>1. Ask if pt. have allergies.</li> <li>2. Notify physician if signs of severe reactions occur.</li> <li>3. Inform pt. to finish all the series of shots.</li> <li>4. Teach pt. that the vaccine does not contain live virus, works by producing immunity that prevent from getting infection.</li> </ol>
Hepatitis B Immune Globulin	Specific immune globulin	Treatment for hepatitis B, liver transplant		Y N		N/V/D, back pain, rash, Pain/swelling/ redness at the injection site, vision problems.	<ol style="list-style-type: none"> <li>1. Notify physician if signs of severe reactions occur.</li> <li>2. Ask if pt. have allergies.</li> <li>3. Inform pt. to notify before getting flu shots or other vaccines while receiving the med.</li> <li>4. Inform pt. that he/she may need to have a second dose of med 1 month after the 1<sup>st</sup> dose.</li> </ol>
				Y N			<ol style="list-style-type: none"> <li>1.</li> <li>2.</li> <li>3.</li> <li>4.</li> </ol>