

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Oxytocic hormone	-induce or strengthen uterine contractions or control bleeding after childbirth	10-40 units, do not exceed 40 units IV	Y N	10 u/1,000ml	Confusion, Convulsions, Difficulty breathing, hives, headache	1. Monitor uterine activity 2. Monitor fetal heart rate 3. Assess fetal presentation and station before administering 4. Weigh pads
Magnesium Sulfate	Tocolytic	Prevent/treat seizures in women with preeclampsia or eclampsia	4-6g IVPB	Y N	Must be diluted to a concentration of 205 or less prior to IV infusion Rate of admin. should be slow and cautious	Breathing difficulties, poor reflexes, confusion, weakness, flushing, sweating	1. Resp. rate should be at least 16 breaths per minute before administering 2. Urinary output should be at least 30ml per hour over 4 hours 3. Check patellar reflexes 4. Loss of DTR=Stop the mag
Meperidine	Opiate analgesics	Treat/relieve severe pain	PO 50-150 mg	Y N	Push over 3-5 minutes (give slowly)	Light headedness, dizziness, weakness, headache, mood changes	1. Do not give if pt. has taken MAOIs in the past 14 days 2. May cause life threatening breathing problems especially during 24-72 hours 3. Don't take with St. John's Wort 4. Monitor closely for signs of sedation
Promethazine	Phenothiazine	Used to relax and sedate	6.25-12.5mg	Y	25mg/ml	Dry mouth, drowsiness,	1. Monitor respirations 2. Assess dizziness/drowsiness, may affect

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Newborn Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: _____

		patients	q 8hrs	N	Administer slowly	nightmares, dizziness, tinnitus, blurred vision	patient's gait and balance. 3. Give IM injections deep into muscle 4. Do not administer intra-arterially!
		Shorten duration of labor					
Calcium Gluconate	Antidotes/ calcium salts	Antidote for magnesium sulfate toxicity	500-2000mg	Y N	Should not be admin. at a rate exceeding 200mg/min	Nausea, vomiting, decreased appetite, constipation, dry mouth	1. Assess for cutaneous burning sensation 2. Monitor HR and ECG 3. Monitor for Chvostek's and Trousseau's signs 4. Monitor lab values

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	Vitamin K	Used to prevent bleeding	1-3 mg IM/IV/ subq	Y N	50ml over 60 minutes should not exceed 1mg/min	Injection site swelling, redness, soreness. Flushing, taste changes, dizziness, tachycardia	2. Infuse slowly 2. Caution in pt.'s with renal impairment 3. Tell pt. to report chest pain 4. Monitor for breathing difficulty
Erythromycin Ophthalmic	Macrolide antibiotics	Treat/prevent eye infections		Y	n/a	Burning, stinging,	1. Pull lower eyelid down gently and squeeze s half inch ribbon into the sac

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: _____

Ointment		of newborn babies	1 cm	N		swelling of face/lips/tongue, redness, allergic reactions	2. Admin. within one hour after birth 3. Obtain informed consent by baby's parents/guardians 4. Monitor for adverse reactions
Engerix B	Vaccine	Immunization against HepB	4 total doses 40mcg	Y N	n/a	Pain/soreness/swelling at injection site. Fever, sore throat, nausea	1. Administer intramuscularly 2. Deltoid for children and adults 3. Under year 1 admin. in anterolateral aspect of the thigh 4. Admin. subq in patients with bleeding disorders
Hepatitis B Immune Globulin	Disease specific immunoglobulin	Immunization against HepB	0.5ml IM	Y N	n/a	Injection site tenderness and redness, anaphylaxis	1. Do not shake 2. monitor for adverse effects 3. Admin. within 12 hours after birth 4. Obtain consent from parent/guardian
				Y N			1. 2. 3. 4.