

Maternal Medication Worksheet– Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)		Circle IVF Type		Rationale for IVF		Lab Values to Assess Related to IVF		Contraindications/Complications	
		Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>							
Student Name: Joshua Rivera		Unit: OB SIM		Patient Initials:		Date: June 6, 2022		Allergies:	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route, & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects		Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)	
Oxytocin	Oxytocic Hormone/ Agent	Hemorrhage Induction Incomplete Abortion	10u/mL IM 10-40 IV			Bleeding, Blurred Vision, Arrhythmias, Fetal death		<ol style="list-style-type: none"> 1. Monitor Intrauterine Pressure. 2. Monitor Fetal Heart Rate. 3. D/C immediately with hyperactivity or fetal distress. 4. If hyperactivity/fetal distress occur, administer oxygen, and put mom in left lateral position. 	
Magnesium Sulfate	Antidysrhythmic	Pre-eclampsia Eclampsia Toxemia	40mg/ml 80mg/ml INJ 1g/100ml 2g/100ml IV			Circulatory Collapse, Hypotension, Respiratory Paralysis, Pulmonary Edema		<ol style="list-style-type: none"> 1. Do not start, stop, or change dose without checking with doctor. 2. Monitor renal function, BP, RR, and deep tendon reflex. 3. Use as last resort, positive evidence of human fetal risk. 4. Safe for use while breastfeeding. 	
Meperidine	Opioid Analgesic	Moderate - Severe Pain	Tablets or Syrup q3-4hr			Seizures, Decreased libido, Agitation, Rash/Hives		<ol style="list-style-type: none"> 1. Can be taken with or without food. 2. Don't chew, break, or crush tablets. 3. Don't stop without talking to your doctor. 4. Use dosage spoon or cup to measure correct dose if syrup. 	
Promethazine	Phenothiazine	Nausea Vomiting Antihistamine	PO			Confusion, Double Vision, Jaundice, Insomnia		<ol style="list-style-type: none"> 1. Taken at bedtime or before meals. 2. How often/dose is determined off condition being treated. 3. Call Dr. if symptoms don't improve or worsen. 4. Do not take extra to make up for a missed dose. 	
Student Name: Joshua Rivera		Unit: OB SIM		Patient Initials:		Date: June 6, 2022		Allergies:	

Maternal Medication Worksheet– Current Medications & PRN for Last 24 Hours

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route, & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Calcium Gluconate	Calcium Salt Antidote	Hypocalcemia Tetany Calcium Supplement	100mg/ml INJ 50,500,650mg Tablets 500mg Capsule			Hypotension, Flatulence, Polyuria, Bradycardia	<ol style="list-style-type: none"> 1. Severe interaction with ceftriaxone. 2. Enters human milk, use caution if breastfeeding. 3. If PO, take with food. 4. Do not double dose if a dose is missed.
Phytonadione	Vitamins (K) Anticoagulant reversal agent	Deficiency Bleeding/Clotting Problems	_____			Cyanotic Lips, Dyspnea, Bradycardia, Jaundice	<ol style="list-style-type: none"> 1. Never use in large amounts, or for longer than recommended by doctor. 2. Store at room temperature away from moisture and heat, protect from light. 3. May need frequent blood tests. 4. Call doctor for instructions if dose is missed.
Erythromycin Ophthalmic Ointment	Macrolide Antibiotic	Prevent/Treat Eye Infections	_____			Vision Changes, Facial Swelling, Itching, Burning	<ol style="list-style-type: none"> 1. Medication is only for use in the eye. 2. Wash hands before and after every use. 3. Try to avoid touching tip of tube to your eye, fingertips, or any other surface. 4. Use at regular intervals, exactly as prescribed for the full course.
Engerix B	Viral Vaccine	Hepatitis B Prevention	INJ			Light-Headed, Fever, Headache, Myoclonic	<ol style="list-style-type: none"> 1. Helps develop immunity, does not treat active infection. 2. Contact doctor if you miss a booster dose or get behind. 3. Receive all doses or you may not be fully protected. 4. Keep track of all side effects you have after receiving to report to Dr.
Hepatitis B Immune Globulin	Specific Immune Globulin	Hepatitis B Exposure Reoccurrence Preventative	_____			Tremors, Joint Pain, Vomiting, Diarrhea	<ol style="list-style-type: none"> 1. Provides immediate, short-term protection. 2. Observe for adverse effects 15 minutes after injection. 3. Give after stabilization of infant, preferably within 12hr after birth. 4. Report adverse effects.