

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
Click here to enter text.	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>	Click here to enter text.	Click here to enter text.	Click here to enter text.

<b>Student Name:</b> Mariah Mata	<b>Unit:</b> SIM	<b>Patient Initials:</b> Click here to enter text.	<b>Date:</b> 11/1/2022	<b>Allergies:</b> Click here to enter text.
-------------------------------------	---------------------	---	---------------------------	--

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Hydroxyzine pamoate	Antihistamine	Anxiety	25mg PO PRN q6	<div style="border: 1px solid black; width: 100px; height: 20px; margin-bottom: 5px;"></div> Click here to enter text.		Dizziness, fainting, convulsions, headache	1. Can cause a serious heart problems especially if used at the same time as other medications. Tell your doctor about all medications taken 2. medication may impair your thinking or reaction so be careful if you drive or do anything that requires you to be alert 3. Do not drink alcohol while taking this medication 4. You should not take this medicine for longer than 4 months
lorazepam	Benzodiazepine	Alcohol withdrawal / anxiety	1-2 mg PO PRN q4 hours	<div style="border: 1px solid black; width: 100px; height: 20px; margin-bottom: 5px;"></div> Click here to enter text.		dizziness, drowsiness, weakness, feeling unsteady	1. Can slow or stop your breathing, especially if you have recently used an opioid medication, alcohol, or other drugs that can slow your breathing. 2. misuse of this medication can cause addiction, overdose, or death 3. avoid drinking alcohol 4. you should not use ativan if you have

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Mariah Mata		Unit: SIM	Patient Initials: Click here to enter text.		Date: 11/1/2022	Allergies: Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							narrow angle glaucoma, severe respiratory insufficiency or, myasthenia gravis
haldol	antipsychotic	for psychotic symptoms/agitation	10 mg IM PRN q8	<input type="text"/> Click here to enter text.		Drowsiness, headache, dizziness, spinning sensation, feeling restless or anxious	<ol style="list-style-type: none"> <li>1. Haldol is not approved for use in older adults with dementia related psychosis</li> <li>2. Do not take medication if you have Parkinson's disease</li> <li>3. Tell doctor if you have ever had long QT syndrome (in you or a family member)</li> <li>4. Never take more medication than prescribed, it can cause serious heart rhythm disorder or sudden death.</li> </ol>
diphenhydramine	antihistamine	management of cholinergic effects of antipsychotic	50 mg IM q8 PRN psychotic symptoms	<input type="text"/> yes	N/A	dizziness, drowsiness, loss of coordination, dry mouth, nose, or throat, constipation, upset stomach	<ol style="list-style-type: none"> <li>1. Use caution driving, operating machinery, or performing other hazardous activities. Medication may cause dizziness or drowsiness</li> <li>2. Use alcohol cautiously. Alcohol may increase drowsiness and dizziness</li> <li>3. store medication at room temperature away from moisture and heat</li> <li>4. If does is missed wait till it is time to take medication again. Do not take extra medication to make up for missed dose</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Mariah Mata		Unit: SIM	Patient Initials: <a href="#">Click here to enter text.</a>	Date: 11/1/2022	<b>Allergies:</b> <a href="#">Click here to enter text.</a>		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
acetaminophen	Analgesic/antipyretic	Mild-moderate pain	500 mg PO q6 for pain of 1-4 on pain scale	<input type="text"/> <a href="#">Click here to enter text.</a>		stomach pain, loss of appetite, tiredness and itching	<ol style="list-style-type: none"> <li>1. Stop taking this medicine and call your doctor right away if you have skin redness or a rash that spreads and causes blistering and peeling</li> <li>2. Be sure to tell your doctor if you are on a blood thinner.</li> <li>3. . Avoid using other medicines that may contain acetaminophen</li> <li>4. An overdose of acetaminophen can damage your liver or cause death.</li> </ol>
olanzapine	antipsychotic	Agitation/psychotic behavior	5 mg PO q6 for agitation	<input type="text"/> <a href="#">Click here to enter text.</a>		dry mouth, tremors, changes in personality, weight gain, increased appetite	<ol style="list-style-type: none"> <li>1. Medication can cause high blood sugar</li> <li>2. Educate patient to stay hydrated. patients can easily get overheated and dehydrated when taking the medication</li> <li>3. Long term use of this medication can cause a serious movement disorder.</li> <li>4. Medication can be taken with or without food</li> </ol>
ibuprofen	NSAID	Mild-moderate pain	400 mg PO q6 for pain of 1-4 on pain scale	<input type="text"/> <a href="#">Click here to enter text.</a>		Changes in vision, swelling or rapid weight gain, N/V, bleeding,	<ol style="list-style-type: none"> <li>1. can increase your risk of fatal heart attack or stroke. Do not use this medicine just before or after heart bypass surgery</li> <li>2. do not take more than your recommended dose an ibuprofen</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

<b>Student Name:</b> Mariah Mata		<b>Unit:</b> SIM	<b>Patient Initials:</b> Click here to enter text.		<b>Date:</b> 11/1/2022	<b>Allergies:</b> Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
						dizziness, headache	overdose can damage your stomach or intestines.  3. you should not take this medication if you have ever had an asthma attack or severe allergic reaction after taking NSAID  4. The maximum amount of medication per day for adults is 3200mg
risperdal	antipsychotic	Psychotic behavior/bipolar mania	1 mg tablet PO BID	<input style="width: 100px; height: 20px;" type="text"/> Click here to enter text.		Headache, dizziness, drowsiness, feeling tired, tremors, twitching or uncontrollable muscle movements, agitation, anxiety	1. medication is not approved for use in older adults with dementia related psychosis  2. while taking medication you may be more sensitive to temperature extremes such as hot or cold conditions. Avoid getting too cold or becoming overheated or dehydrated  3. stop using and call your doctor at once if you have fever, stiff muscles, confusion, sweating, fast or uneven heartbeats, restless muscle movements in your face or neck.  4. If you become pregnant while taking, do not stop taking it without your doctor's advice

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Mariah Mata		Unit: SIM	Patient Initials: <a href="#">Click here to enter text.</a>	Date: 11/1/2022	<b>Allergies:</b> <a href="#">Click here to enter text.</a>		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Nicotine patch	Stimulant	Nicotine replacement	21 mg transdermal patch daily	<input style="width: 100px; height: 20px;" type="text"/> <a href="#">Click here to enter text.</a>		HTN, mood changes, chest pain or pressure, dizziness, numbness/tingling feeling, cold sweats, change in hearing	<ol style="list-style-type: none"> <li>1. Make sure previous patch is removed before placing the new one</li> <li>2. make sure to throw away old patch in black container in the med room</li> <li>3. Take patch off before having an MRI</li> <li>4. watch blood sugar closely can cause high blood glucose</li> </ol>
ziprasidone	antipsychotic	Psychotic behavior/bipolar disorder/mania	40 mg PO BID	<input style="width: 100px; height: 20px;" type="text"/> <a href="#">Click here to enter text.</a>		N/V, trouble swallowing, weight gain, vision problems, headache, dizziness, drowsiness, weakness	<ol style="list-style-type: none"> <li>1. do not take medication if you have a heart rhythm disorder, Long QT syndrome, uncontrolled heart failure, or if you have recently had a heart attack</li> <li>2. stop taking medication and call doctor right away if you have sudden dizziness, fast or pounding heartbeats, fluttering in your chest, SOB</li> <li>3. medication is not approved for use in older adults with dementia related psychosis</li> <li>4. It may take several weeks before symptoms improve. Do not stop taking</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Mariah Mata		Unit: SIM	Patient Initials: <a href="#">Click here to enter text.</a>	Date: 11/1/2022	<b>Allergies:</b> <a href="#">Click here to enter text.</a>		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							medication suddenly even if you feel fine
prazosin	Alpha-blocker	Nightmares	1 mg PO QHS	<input type="text"/>  <a href="#">Click here to enter text.</a>		Feeling weak or tired, nausea, dizziness, drowsiness, light headed feeling	<ol style="list-style-type: none"> <li>1. medication can affect your pupils. If you have cataract surgery, tell your surgeon ahead of time that you are taking this medication</li> <li>2. not approved for use by anyone younger than 18 years old</li> <li>3. medication lowers blood pressure and may cause dizziness or fainting, especially when you first start taking it or whenever dose is changes</li> <li>4. blood pressure will need to be checked often</li> </ol>
Nicotine inhaler	Stimulant	Nicotine replacement	10 mg cartridge for inhalation Q2	<input type="text"/>  <a href="#">Click here to enter text.</a>		Upset stomach, diarrhea, drooling, shakiness, feeling tired or weak, cold sweats, blurred eyesight, changes in hearing	<ol style="list-style-type: none"> <li>1. Do not use more than told. Unsafe side effects may happen</li> <li>2. tell your doctor if you are taking another drug to help stop smoking</li> <li>3. IF you have high blood sugar, you will need to watch your blood sugar closely</li> <li>4. Notify the doctor if you have not been able to quit smoking after using nicotine inhaler for as long as it says to use it.</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

<b>Student Name:</b> Mariah Mata		<b>Unit:</b> SIM	<b>Patient Initials:</b> Click here to enter text.		<b>Date:</b> 11/1/2022	<b>Allergies:</b> Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	<input type="text"/> Click here to enter text.	<input type="text"/> here to enter text.	Click here to enter text.	<ol style="list-style-type: none"> <li>1. Click here to enter text.</li> <li>2. Click here to enter text.</li> <li>3. Click here to enter text.</li> <li>4. Click here to enter text.</li> </ol>
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	<input type="text"/> Click here to enter text.	<input type="text"/> here to enter text.	Click here to enter text.	<ol style="list-style-type: none"> <li>1. Click here to enter text.</li> <li>2. Click here to enter text.</li> <li>3. Click here to enter text.</li> <li>4. Click here to enter text.</li> </ol>