

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
Ondansetron	Route: oral		Is used to prevent nausea and vomiting that may be caused by surgery, cancer chemotherapy, or radiation treatment.	<ol style="list-style-type: none"> 1. Side effects may include constipation or diarrhea 2. Instruct patient to report symptoms of serious cardiac arrhythmias (change in heart rate, lightheadedness, syncope) 3. Avoid activities requiring mental alertness until drug effects are known. Drug may cause fatigue or malaise 4. Instruct patient to let oral film dissolve and not to chew or swallow it
Metoprolol	Route: oral		Metoprolol is a beta-blocker used to treat chest pain or hypertension.	<ol style="list-style-type: none"> 1. Avoid activities requiring mental alertness until drug effects are known. Drug may cause dizziness. 2. Instruct patient to inform physician before any kind of surgery that drug is being used 3. Teach patient not to stop the drug suddenly because it may cause hypertension, angina, or myocardial infarction.

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

				<ol style="list-style-type: none"> 4. Advise patient to report symptoms of new or worsening heart failure
Levofloxacin	Route: oral		<p>Levofloxacin is an antibiotic that may be used to treat different types of bacterial infections. It is usually only used to treat bacterial infections that cannot be treated with safer antibiotics.</p>	<ol style="list-style-type: none"> 1. Immediately report symptoms of tendonitis, tendon ruptures, or peripheral neuropathy. 2. Use sunscreen/wear protective clothing due to potential for photosensitivity 3. Avoid activities requiring mental alertness until drug effects are known. Drug may cause dizziness, lightheadedness. 4. Immediately report symptoms of an aortic aneurysm (sudden chest, stomach or back pain)
Vancomycin	Route: oral		<p>Vancomycin is an antibiotic used to fight bacteria. It is most commonly used to treat C. Diff.</p>	<ol style="list-style-type: none"> 1. Use medicine for the full prescribed length of time, even if symptoms quickly improve. Skipping doses can increase risk of infection that is resistant to medication. 2. May cause low potassium levels 3. Report hearing loss or ringing in ears

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

				<ol style="list-style-type: none"> Report kidney problems such as rapid weight gain, pain on side or lower back, little or no urination
Prednisone	Route: oral		<p>Prednisone is used to decrease inflammation/ It can be used to treat allergic disorders, skin conditions, asthma, COPD and many more conditions.</p>	<ol style="list-style-type: none"> Drug may cause N,V,D Advise patient against sudden discontinuation of drug Avoid alcohol and cigarettes, it may increase risk for osteoporosis Advise patient to monitor blood glucose due to drug-induced hyperglycemia

Allergies:

Primary IV fluid and rate:

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Medication reference:

Merative (2022). Micromedex Drug Reference (Version 3.0.0) [Mobile App]. App Store.

<https://apps.apple.com/us/app/micromedex-drug-reference/id666520138>

Drugsite Trust (2022). Drugs.com Medication Guide (Version 2.10.9) [Mobile App]. App Store.

<https://apps.apple.com/us/app/drugs-com-medication-guide/id599471042>