

Student Name: Jamie Dominguez

Unit: Pedi AM

Pt. Initials: Appendicitis

Date: 1/25/23

Pediatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: Unknown

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
D5 1/2 NS @ 116 mL/hr	Isotonic/ Hypotonic / Hypertonic	As maintenance to push fluid from the intravascular space into the cell	Sodium and Blood Glucose	fever, infections at the site of injection, blood clot or inflammation surround the site of injection, leakage of fluid into the surrounding tissues (extravasation),

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Therapeutic Range?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				Is med in therapeutic range?			
				If not, why?			
Meropenem	Carbapenem Antiinfective	interferes with Cell-wall Replication of susceptible organism Preventative for Appendicitis	1 gram IV every 8 hours	>50 kg (1 gram IV every 8 hours) Yes in range	Reconstitute with 20 ml of NS and administer over 15 to 30 minutes in pump	Diarrhea Glossitis Rash Vomiting	1. Report any itching, rash, shortness of breath 2. Promptly report diarrhea or bloody stools 3. May interfere with mental alertness 4. Report any burning, pain or stinging at injection site.
							1. 2. 3. 4.
							1. 2. 3. 4.
							1. 2. 3. 4.
							1. 2. 3. 4.

Student Name: Jamie Dominguez

Unit: Pedi AM

Pt. Initials: Atopic Dermatitis

Date: 1/25/23

Pediatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: Unknown

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Therapeutic Range?	IVP – List solution to dilute and rate to push. IVPB – concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				Is med in therapeutic range?			
				If not, why ?			
Pimecrolimus	Misc. topical agent	Decrease redness, swelling, pain	Apply thin layer topically twice daily and rub in.	Children > 2 yrs of age apply thin layer topically twice daily and rub in. YES		Burning	<ol style="list-style-type: none"> 1. Advise patient to avoid contact between medication and eyes 2. Instruct patient to discontinue use of product when condition clears. 3. Assess skin conditions 4. Assess for skin infections
Levocetirizine	Antihistamine	Decreases allergic response by blocking histamine therefore reduces itching.	1.25 mg by mouth once daily in the evening	1.25 mg by mouth once daily in the evening YES		Drowsiness fatigue dizziness hepatitis dry mouth increased LFT's urinary retention	<ol style="list-style-type: none"> 1. Notify MD if patient has confusion, sedation, hypotension 2. To avoid taking any other CNS depressants 3. Assess respiratory status 4. Monitor Liver Function Test
							<ol style="list-style-type: none"> 1. 2. 3. 4.
							<ol style="list-style-type: none"> 1. 2. 3. 4.
							<ol style="list-style-type: none"> 1. 2. 3. 4.