



<p>7. Pain & Discomfort Management: List 2 Developmentally Appropriate Non-Pharmacologic Interventions Related to Pain & Discomfort for This Patient.</p> <ol style="list-style-type: none"> 1. Reposition pt. to relieve ICP 2. Pacifier use <p>*List All Pain/Discomfort Medication on the Medication Worksheet</p>	<p>8. Calculate the Maintenance Fluid Requirement (Show Your Work): Patient Wt: 8 kg</p> <p>$8 * 100 = 800 \text{ mL}$ $800 \text{ mL} / 24 \text{ hr}$</p> <p>Calculated Fluid Requirement: 33.3 ml/hr</p> <p>Actual Pt MIVF Rate: _____ ml/hr</p> <p>Is There a Significant Discrepancy?</p> <p>Why?</p>	<p>9. Calculate the Minimum Acceptable Urine Output Requirement (Show Your Work): $8 \text{ kg} * 1 \text{ mL/kg/hr}$</p> <p>Calculated Min. Urine Output: 8 ml/hr</p> <p>Actual Pt Urine Output: _____ ml/hr</p>
<p>10. Growth & Development: List the Developmental Stage of Your Patient For Each Theorist Below and Document 2 OBSERVED Developmental Behaviors for Each Theorist. If Developmentally Delayed, Identify the Stage You Would Classify the Patient: Patient age: 8 Months</p> <p>Erickson Stage: trust vs mistrust</p> <ol style="list-style-type: none"> 1. Comforted with physical touch 2. relies on caregivers <p>Piaget Stage: Sensorimotor</p> <ol style="list-style-type: none"> 1. sits up 		

Student Name [Hans Burda](#)

	2. crawls
--	------------------

<p>11. Focused Nursing Diagnosis: - Risk for injury</p>	<p>15. Nursing Interventions related to the Nursing Diagnosis in #11: 1. Measure newborn’s head circumference</p> <p>Evidenced Based Practice: - allows for adequate ICP management</p> <p>3. Monitor LOC</p> <p>Evidenced Based Practice: - Checking pupils for equality and reaction</p> <p>3. Monitor neurological status</p> <p>Evidenced Based Practice: - Observe shrill cry, lethargy or irritability</p>	<p>16. Patient/Caregiver Teaching: 1. Reinforce doctor’s education about the surgical procedure 2. Educate on proper care of shunt at home</p> <p>3. Report S/S of infection</p>
<p>12. Related to (r/t): - related to increased ICP</p>		<p>17. Discharge Planning/Community Resources: 1. Pharmacy</p> <p>2. Medical library</p> <p>3. Telehealth</p>
<p>13. As evidenced by (aeb): - as evidenced by bulging anterior fontanelles</p>		
<p>14. Desired patient outcome: - prevention of injury and management of ICP</p>		