

Pediatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: NKDA

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
N/A	Isotonic/ Hypotonic/ Hypertonic	N/A	N/A	N/A

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Therapeutic Range?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions  (Precautions/Contraindications, Etc.)
				Is med in therapeutic range?			
				If not, why?			
<b>Pediatric Multivitamin Drops with Iron (Polyvisol)</b>	Iron products; Vitamin and mineral combinations	Used to help growth and good health	10mcg/10mL PO Every 12 hours	<b>0.5 - 1mL (or as indicated by the individual) They are being given more, but they also need more iron than a full term baby</b>	N/A	Black, tarry or bloody stools Fever Upset stomach or throwing up	1. Nurses need to know when the last dose was given because overdose can lead to toxicity. 2. Teach parents/nurses monitor for bloody or black stools. 3. Baby may have an upset stomach so note how much they are eating and if their abdomen is distended. 4. If baby throws up, note what it looks like, if it looks like coffee grounds or blood it could be from drops.
							1. 2. 3. 4.
							1. 2. 3. 4.