



<p>7. Pain & Discomfort Management: List 2 Developmentally Appropriate Non-Pharmacologic Interventions Related to Pain & Discomfort for This Patient.</p> <ol style="list-style-type: none"> 1. Repositioning 2. Provide calm environment <p>*List All Pain/Discomfort Medication on the Medication Worksheet</p>	<p>8. Calculate the Maintenance Fluid Requirement (Show Your Work): Patient Wt: 64.4 kg</p> <p>10 * 100 = 1000mL 10 * 50 = 500mL 44.4 * 20 = 888mL 2388mL/24hr</p> <p>Calculated Fluid Requirement: 99.5 ml/hr</p> <p>Actual Pt MIVF Rate: 0 ml/hr</p> <p>Is There a Significant Discrepancy?</p> <p>Why?</p>	<p>9. Calculate the Minimum Acceptable Urine Output Requirement (Show Your Work): 64.4kg * 0.5mL/kg/hr</p> <p>Calculated Min. Urine Output: 32.2 ml/hr</p> <p>Actual Pt Urine Output: <u>not measured</u></p>
	<p>10. Growth & Development: List the Developmental Stage of Your Patient For Each Theorist Below and Document 2 OBSERVED Developmental Behaviors for Each Theorist. If Developmentally Delayed, Identify the Stage You Would Classify the Patient: Patient age: 9 years</p> <p>Erickson Stage: Industry vs Inferiority</p> <ol style="list-style-type: none"> 1. Admits to playing sports 2. Does schoolwork <p>Piaget Stage: Concrete operational</p> <ol style="list-style-type: none"> 1. Able to use laptop 2. Able to navigate through halls 	

<p>11. Focused Nursing Diagnosis: - Risk for bleeding</p>	<p>15. Nursing Interventions related to the Nursing Diagnosis in #11: 1. Review lab results</p> <p>Evidenced Based Practice: - monitoring PLT, PT/INR levels to determine effectiveness of therapy.</p> <p>3. Prevent injury</p>	<p>16. Patient/Caregiver Teaching: 1. Refrain from contact sports 2. Report S/S of bleeding</p> <p>3. Report S/S of infection</p>
<p>12. Related to (r/t): - related to decreased PLT count</p>	<p>Evidenced Based Practice: - Provide safe surroundings and put call light within patient reach reduces the risk for injury.</p> <p>3. Prevent infections</p> <p>Evidenced Based Practice: - Perform hand hygiene and clean stethoscope before each use to reduce the spread of infection</p>	
<p>13. As evidenced by (aeb): - as evidenced by bruising and CBC results</p>		<p>17. Discharge Planning/Community Resources: 1. Pharmacy</p> <p>2. Medical library</p> <p>3. Telehealth</p>
<p>14. Desired patient outcome: - prevention of bleeding</p>		