

Student Name: Joel Castillo

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: _____

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Oxytocic Nimines	to help start or continue labor + controlling bleeding after del.	0.5-1 mu IVP	Y N	IVP- q 15-60 min until contraction pattern reached	confusion renal/urinary diff. breathing tachycardia MA, hiccups	1. monitor fetal heart rate 2. monitor I/O water intake/output common 3. contraindicated w/ fetal distress, prematurity 4. advise pt. to expect contractions similar to menstrual cramps
Magnesium Sulfate	antidysrhythmic (electrolytes)	prevent seizures in preeclampsia	100mg/kg in 4-8 hrs oral	Y N	N/A	hypotension pulmonary edema reflexes hypertension flushing dizziness	1. monitor respiratory status, deep tendon reflexes 2. monitor electrolyte levels for hypermagnesemia 3. calcium gluconate is the antidote 4. contraindicated w/ renal failure pts.
Meperidine	phenylpiperidine	treatment of moderate to severe pain	50-100 mg orally q3-4hrs	Y N	N/A	lightheadedness MA, agitation constipation blurred vision	1. call before fall to ↓ risk of falling 2. ↑ water intake and promote high-fiber diet 3. narcosis is the antidote 4. monitor bp prior to administration can cause resp. depression
Promethazine	phenothiazines	used to relieve symptoms of allergic rxn	25 mg twice daily orally	Y N	N/A	dry mouth dizziness disturbances diff. falling drowsiness	1. call before fall when needing to get up 2. contraindicated w/ low BP 3. can be taken w/ or w/o food 4. report mental status, fever, diff. breathing
Calcium Gluconate	antidotes	manage hypocalcemia, cardiac arrest, & cardiotoxicity due to hyperkalemia/hypermagnesemia	oral 500-2000mg 2-4x a day	Y N	N/A	chalk taste nausea constipation nausea, tingling heavy feeling	1. ↑ intake of milk, dark green vegetables 2. avoid sun-rin food inhibits calcium absorption 3. frequent labs to assess electrolyte imbalance 4. report S/S of hyp/hyper calcemia

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	vitamin K fat-soluble	prevent bleeding in people w/ blood clotting problems or too little vit. K in the body	2.5-10mg orally daily	Y N	N/A	nausea, diff. breathing, weakness, lightheadedness itching, chest pain	2. report S/S of distress, lightheadedness, S/S of hypotension 3. contraindicated w/ severe renal failure/hitrate 3. frequent PT/INR lab tests for clotting time 4. avoid ↑ intake of vit. K when on drug regimen
Erythromycin Ophthalmic Ointment	macrolide antibiotic	treat bacterial infections of the eye	once at birth	Y N	N/A	eye redness mild irritation discomfort itching	1. ensure proper administration of good hygiene, antibiotic 2. call before fall since and will be on eye pt. will be limited for a 3. report any pain, swelling, worsen may be S/S of allergic rxn 4. monitor liver fxa tests, can be toxic if used unsafely
Engerix B	vaccines (inactivated, viral)	immunization against infections caused by all known subtypes of Hep. B virus	1m, 1m, 2-dose 0, 1, 6mo, boost	Y N	N/A	pain, soreness, swelling at inj site fever, MA, soreness, itchy head, dizziness	1. contraindicated w/ anaphylaxis to serum/ova 2. report S/S of diff. breathing, cyanosis, swelling of face, throat 3. be aware of previous hep B vaccine dx 4. ensure consent and proper administration
Hepatitis B				Y			1. administer into deltoid muscle or anterolateral of thigh for born absorption

Immune Globulin	IMMUNE globulins	prevent hep. B from occurring again	0.10 mL/kg/dose	N	N/A	<p>1. HIV, infectious mononucleosis, hepatitis, toxoplasmosis, febrile response</p> <p>2. Have epinephrine readily available for adverse rxn</p> <p>3. Do not breast-feed after taking this drug consult w/ physician</p> <p>4. administer w/ calculation of hep B vaccine <small>might impair active humoral response</small></p>
				Y		<p>1.</p> <p>2.</p> <p>3.</p> <p>4.</p>
				N		