

## Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
Allopurinol (Zyloprim)	300 mg PO daily		Treatment of pts Gout	<ol style="list-style-type: none"> <li>1. Report any numbness/tingling arms/hands, legs/feet</li> <li>2. Drink plenty of fluids but watch for changes in voiding (frequency, amount, color, pain during urination, and blood in urine)</li> <li>3. Stop taking this med and call PCP if you develop a skin rash especially if taking amoxicillin</li> <li>4. Can be helpful to take after eating to avoid upset stomach</li> </ol>
Amlodipine (Norvasc)	10 mg PO daily		Treatment of pts hypertension	<ol style="list-style-type: none"> <li>1. Be careful when getting up from a laying or sitting position</li> <li>2. Report any swelling in face hands and feet</li> <li>3. Watch for headaches dizziness and fatigue</li> <li>4. Call PCP if you feel like your heart is racing</li> </ol>
Apixaban (Eliquis)	5 mg PO BID		Used to prevent blood clots since pt has a hx of a-fib	<ol style="list-style-type: none"> <li>1. Report blood/black/tarry stool</li> <li>2. Severe headaches, unusual fatigue, vision changes and slurred speech.</li> <li>3. If you need to go to the dentist make sure to let them know you take this medication and report any gum bleeding.</li> </ol> <p><b>**DO NOT ABRUPTLY STOP</b></p>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

Apixaban (Eliquis) cont.				TAKING THIS MED UNLESS COVERAGE WITH ANOTHER ANTICOAGULANT IS ALSO BEING GIVEN, COMPLETEION OF THERAPY, OR BLEEDING HAS OCCURRED. HIGHER RISK OF HAVING A STROKE DUE TO HAVING A HX OF A-FIB. **
Docusate/Senna (Senokot-S)	50/8.6 mg 1 Tab PO BID		Constipation	1.abdominal cramping 2.Yellow or brown urine 3.Obstruction/fecal impaction 4.GI or rectal bleeding
Duloxetine (Cymbalta)	60 Mg daily		Treatment of pts diabetic neuropathy	1.Check and treat blood sugar as often as needed 2.report urine retention and s/s of UTI 3.Check blood pressure while taking this medication 4. stop taking medication at the first sign of rash with blistering **ADVISE FAMILIES AND CAREGIVERS OF THE NEED FOR CLOSE OBSERVATION AND COMMUNICATION WITH THE PRESCRIBER INCASE OF SUICIDAL THOUGHTS AND BEHAVIORS**
Ferrous Sulfate (Slow FE)	325 mg daily with breakfast		Treatment of pts iron deficiency	1.Constipation/stomach cramps 2.Avoid taking antacids within 2 hrs before or after taking this medication and wait ten mins before laying down. 3.This med can temporarily stain your teeth
Ferrous Sulfate				

## Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

(Slow FE) Cont.				4.stop taking this med and report any s/s of an allergic reaction <b>** Keep out of reach of children. Accidental overdose in children under 6**</b>
Gabapentin	300 mg po morning and night		Treatment of pts neuropathic pain due to diabetes	1.Changes in mood, thoughts or behavior 2.report any severe increase in production with cough (pulmonary edema) 3.muscle pain/joint swelling 4.Treat high blood sugar
Lisinopril	5 mg po daily		Treatment of pts hypertension and heart failure	1.stop taking and report s/s of an allergic reaction 2.report any increase in production when coughing or feel increase of shortness of breath 3. call PCP if you have any swelling especially in face tongue or throat. 4.can cause severe low blood pressure so make sure to ssit or stand up slowly to avoid fainting and dizziness <b>**DISCONTINUE USE AS SOON AS POSSIBLE WHEN PREGNANCY IS DETECTED**</b>
Pravastatin  Pravastatin Cont.	20 mg po daily		Treatment of pts high cholesterol	1.Can sometimes change the way food tastes 2.this medication can cause high blood sugar so to treat it as soon as possible 3.Report muscle pain or weakness accompanied by a fever

## Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

				4. Report sudden changes in voiding habits including pain and increased frequency
				1. 2. 3. 4.

Allergies:

Primary IV fluid and rate:

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Medication reference: Sykscape, Medscape, Medlineplus.gov