

Student Name: Amber Peters

Unit: Pedi - med surg

Pt. Initials: 76

Date: 4-4-23

Allergies: NKDA

Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Therapeutic Range?		Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
				is med in therapeutic range?	If not, why?			
Famotidine	ulcer drugs	used prophylactically to prevent ulcers & excess gastric juices	60mg BID	1mg/kg q 12hrs 0.1mg/kg being used properly lactated	IVP 0.6mL = 6mg of 10mg/mL			
Sodium chloride 0.9% Flush	Minerals & electrolytes	used to flush central line sites	10mL every 12 hours (BID)		Intracatheter			
dexamethasone IV syringe (60ir med)	non narcotic analgesic	non narcotic pain control	15mg/kg x 2.2kg	yes	IVPB 35mL = 30mg of 10mg/mL 140mL/hr infused over 15min			
TPN Pediatric Cyclic	minerals, electrolytes, nutrients, multivitamins	to provide nutrition while resting stomach	IV 60mL/hr continuous	0.5-0.7g/kg used as maintenance fluid	Central line infusion 1/2 micron filter			

  

Generic Name	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Famotidine	<ol style="list-style-type: none"> <li>notify nurse of blood in stool</li> <li>may cause fast heartbeat</li> <li>let nurse know/weight</li> <li>for signs depression</li> </ol>
Sodium chloride 0.9% Flush	<ol style="list-style-type: none"> <li>Fluid overload</li> <li>air embolus</li> <li>use sterile technique when accessing / flush</li> </ol>
dexamethasone IV syringe (60ir med)	<ol style="list-style-type: none"> <li>1mg cause nausea &amp; vomiting</li> <li>Do not use more than 4,000 mg/day</li> <li>Monitor hepatic labs.</li> </ol>
TPN Pediatric Cyclic	<ol style="list-style-type: none"> <li>microclots</li> <li>electrolyte imbalance</li> <li>Hyperglycemia</li> <li>Hyperkalemia</li> <li>use sterile technique to avoid central line infection</li> </ol>

Student Name: Amber Peters

Unit: Pedi med surg

Pt. Initials: SD

Date: 4-4-23

Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: Loratadine, Amoxicillin

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Therapeutic Range?		IVP - List solution to dilute and rate to push. IVPB - concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				Is med in therapeutic range? If not, why?				
Cefazolin	cephalosporins	prophylactic for infection after surgery	1,000 mg IVP every 8 hours	25 mg / 50 mg / kg a little above the range due to surgery	IVP 1,000 mg in 10 mL sterile water push over 3-5 min	may cause nausea & diarrhea	1. make sure patient not allergic to 2. monitor prolonged nausea 3. diarrhea 4. let nurse know mucous in stool	
ketorolac	Anti-Pneumatic	IVP for postoperative pain	IVB 15 mg / 15 mg / kg every 6 hours	5 mg / kg yes	IVP Do 15 mL = 23.4 mg of 30 mg / mL	can cause ulcer	1. Do not use concurrently with NSAIDs 2. Can cause ulcer 3. Alert nurse blood in stool 4. Do not use with bleeding disorder	
acetaminophen		mild to moderate pain control	650 mg PO Q4 PRN	300 mg / 10-15 mg / kg yes	325 each tablet PO	in excess can cause liver injury	1. Do not take more than 4,000 mg / day 2. 3. 4.	