

Student Name: \_\_\_\_\_Chelsie Callesen\_\_\_\_\_

Unit:\_\_\_SIM\_\_\_\_\_

Pt. Initials: \_\_\_\_\_

Date: 4/11/2023

**Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	Nonapeptide, pleiotropic hormone	Used to induce labor, strengthen uterine contractions, or to control bleeding after childbirth		Y N		Confusion, seizures, difficulty in breathing, fast or irregular heartbeat, headache, hives, pelvic or abdominal pain	<ol style="list-style-type: none"> <li>1. Monitor contractions and other vital signs closely while administering</li> <li>2. During labor, monitor baby's heart rate closely on the fetal heart monitor</li> <li>3. If uterine hyperactivity occurs, discontinue immediately.</li> <li>4. Monitor for water intoxication, restricting fluid intake may be warranted.</li> </ol>
Magnesium Sulfate	Sulfate, Anti-convulsant	Used to prevent seizures due to worsening preeclampsia, to slow or stop preterm labor, and to prevent injuries to a preterm baby's brain		Y N		Hypocalcemia, skeletal demineralization, osteopenia, and other skeletal adverse effects	<ol style="list-style-type: none"> <li>1. Avoid using for more than 5-7 days for preterm labor (may ↑ risk of hypocalcemia and bone changes in newborn)</li> <li>2. Monitor for changes in pts neuro status</li> <li>3. Avoid continuous use during active labor or within 2 hr. of delivery (potential for magnesium toxicity in newborn)</li> <li>4. Monitor intake and outtake ratios</li> </ol>
Meperidine	Opiate (narcotic) analgesics	Used for moderate to severe pain		Y N		Lightheadedness, headache, agitation,	<ol style="list-style-type: none"> <li>1. Monitor respiratory status (can cause life-threatening breathing problems)</li> <li>2. Use caution in pregnancy: baby may experience life-threatening withdrawal</li> </ol>

**Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

						constipation, uncontrollable shaking of a part of the body, blurred vision, dry mouth	symptoms after birth with regular use during pregnancy 3. Call before getting out of bed and don't get up too quickly; can cause dizziness, lightheadedness, and fainting 4. Take a missed dose as soon as you remember, do not double dose.
Promethazine	Phenothiazines	Used to treat allergic conditions, nausea and vomiting, motion sickness, and sedation		Y N		Drowsiness, headaches, nightmares and feeling dizzy, restless or confused	1. Monitor respiratory status for signs of respiratory depression 2. Can be excreted into breast milk: breastfed infants exposed to this drug should be monitored for sedation, irritability, and excitement 3. If you are having surgery, including dental surgery, tell the doctor or dentist that you are taking promethazine 4. Plan to avoid unnecessary or prolonged exposure to sunlight and to wear protective clothing, sunglasses, and sunscreen. Promethazine may make your skin sensitive to sunlight.
Calcium Gluconate	Calcium Salts	Used to manage hypocalcemia, cardiac arrest, and cardiotoxicity due to hyperkalemia or hypermagnesium		Y N		Nausea, vomiting, decreased appetite, constipation, dry mouth, increased thirst, increased urination, tingling sensations	1. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time(s) each day 2. Do not take other supplements/vitamins unless ordered by your doctor. 3. This medication passes into breast milk. Consult your doctor before breast-feeding. 4. Vitamin D helps with the absorption of calcium. Foods rich in vitamin D include: fortified dairy products, eggs, sardines, cod liver oil, chicken livers, and fatty fish

**Maternal Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	Fat-soluble vitamin	Used to prevent bleeding in people with blood clotting problems or too little vitamin K in the body		Y N		Cyanosis; difficulty swallowing; dizziness, faintness, or lightheadedness; fast, pounding, or irregular heartbeat or pulse; hives, itching, skin rash.	<ol style="list-style-type: none"> <li>1. Never use phytonadione in larger amounts, or for longer than recommended by your doctor</li> <li>2. It may not be safe to breast-feed a baby while you are using this medicine. Ask your doctor about any risks</li> <li>3. May need frequent blood tests</li> <li>4. If you need surgery or dental work, tell the surgeon or dentist you currently use this medicine.</li> </ol>
Erythromycin Ophthalmic Ointment	Macrolide Antibiotics	Used to treat bacterial infections of the eye		Y N		Eye redness or mild eye irritation; severe burning, stinging, or irritation	<ol style="list-style-type: none"> <li>1. Vision may be blurry for a short amount of time after using the eye ointment - this is normal</li> <li>2. It may not be safe to breast-feed while using this medicine. Ask your doctor about any risk</li> <li>3. Use as prescribed, even if your symptoms have improved. Skipping doses can increase the risk of infection that is resistant to medication</li> </ol>

**Newborn Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

							4. Get emergency medical help if you have signs of an allergic reaction: hives; difficult breathing; swelling of your face, lips, tongue, or throat.
Engerix B	Vaccines	Used to help prevent infection from the hepatitis B virus		Y N		Pain, soreness, redness, swelling at the injection site, fever, headache, tiredness, sore throat, nausea, diarrhea, loss of appetite, and dizziness	<ol style="list-style-type: none"> <li>1. You should not receive Engerix-B if you are allergic to yeast</li> <li>2. Engerix-B will not protect against infection with hepatitis A, C, and E, or other viruses that affect the liver</li> <li>3. Be sure to receive all recommended doses or you may not be fully protected against disease</li> <li>4. Cover all cuts carefully. Avoid sharing sharp items such as razors, nail clippers, toothbrushes, and earrings or body rings</li> </ol>
Hepatitis B Immune Globulin	Immune Globulins	Used to provide protection against HBV infection		Y N		Back pain, general feeling of discomfort, headache, muscle aches or pain, pain at the injection site	<ol style="list-style-type: none"> <li>1. This medicine may be given to a baby if the baby's mother has hepatitis B</li> <li>2. Blood tests may be needed to check for unwanted effect</li> <li>3. If using insulin or other medicine for diabetes, you will need to be aware of the symptoms of hypoglycemia (confusion, irritability, double or blurred vision, and in severe cases seizures or loss of consciousness) because this medicine may affect the results of blood sugar tests</li> <li>4. This medicine may cause blood clots. Let doctor know if experiencing chest pain, severe leg pain, SOB, problems with vision or speech</li> </ol>