

Student Name: Miranda Jackson

Unit: _____

Pt. Initials: _____

Date: _____

Maternal Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: _____

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
n/a	Isotonic/ Hypotonic/ Hypertonic	n/a	n/a	n/a

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxytocin	uterine stimulant	labor induction or augmentation post partum hemorrhage	labor: 0.5-2 units/min max: 20 units/min	<input checked="" type="radio"/> Y <input type="radio"/> N	n/a	uterine hyperstimulation uterine tetany uterine rupture abruptio placentae cervical laceration vaginal laceration placental hemorrhage postpartum hemorrhage hypotonia HTN, severe SIADH	1. BBN: not indicated for elective labor (defined as labor initiation w/o medical indications) Contraindicated for fetal malpresentation, uterine cord prolapse, non reabsorbing fetal membranes 2. Avoid uterine pumps, devices @ risk for uterine rupture and acute placental abruption 3. Avoid use in clients receiving vasopressors or prostaglandins like Carboprost 4. Some common side effects are nausea, vomiting, uterine stimulation, hypotension, cardiac arrhythmias and water intoxication
Magnesium Sulfate	anticholinergic	labor suppression	2-6 g IV over 15 to 20 min (loading dose) 1-2g/hr (cont. infusion)	<input checked="" type="radio"/> Y <input type="radio"/> N	always hang w/ isotonic solution	cardiovascular collapse respiratory paralysis hypotension decreased cardiac function pulmonary edema	1. common side effects include depressed reflexes, hypotension, flushing, dizziness, hypocalcemia, and vision changes 2. call before trying to get out of bed 3. monitor serum magnesium levels 4. monitor for mag. toxicity (respiratory depression, chest pain, neuronal inhibition, DTRs, flaccidity, weakness, hyporeflexia)
Meperidine	opioid	Sedation	50-150 mg PO/SC/IM/IV 93-4hr PRN	<input checked="" type="radio"/> Y <input type="radio"/> N	n/a	respiratory depression respiratory arrest dyspnea cardiac arrest syncope seizures bradycardia hypotension	1. BBN: has risk of addiction, abuse and misuse, which can lead to overdose and death. 2. Do not use w/ meprob with 14 days 3. Contraindicated for pts w/ acute and severe asthma due to respiratory depression 4. Call, don't fall!
Promethazine	antiemetic	nausea/vomiting	12.5-25 mg PO 94-4hr PRN	<input checked="" type="radio"/> Y <input type="radio"/> N	n/a	respiratory depression dyspnea seizures thrombocytopenia bradycardia tachycardia NMS	1. Common side effects include drowsiness, sedation, blurred vision, dizziness 2. make sure to call before getting up to prevent falls 3. BBN: contraindicated in pts < 2y/o due to cases of respiratory depression, some fatal 4. BBN: Promethazine injection can cause severe chemical irritation and tissue damage, including burning pain, thrombotic, tissue necrosis and gangrene.
Calcium Gluconate	mmmm	calcium disorders parathyroid disorder: hypoparathyroidism	1-2g IV over PRN	<input checked="" type="radio"/> Y <input type="radio"/> N	n/a	hypercalcemia syncope arrhythmia	1. To be use as an antidote for mag. sulfate toxicity 2. Contraindicated for those that have hypercalcemia 3. monitor electrolyte levels. 4. Report any nausea/vomiting, constipation, or arrhythmias as these are common side effects

Student Name: _____

Unit: _____

Pt. Initials: _____

Date: _____

Newborn Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: _____

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
n/a	Isotonic/ Hypotonic/ Hypertonic	n/a	n/a	n/a

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Phytonadione	Vitamin	Anticoagulant reversal agent	2.5-10mg PO	<input checked="" type="radio"/> Y N	n/a	Warfarin resistance anaphylaxis	<ol style="list-style-type: none"> Common side effects are taste changes w/ SC, IM, or IV use. Flushing, injection site hematoma, and injection site pain. BBN: Some reactions including death have occurred during and immediately after IV infusion even w/ precautions to dilute medication and avoid rapid infusion. Take to your doctor if you have warfarin, Coumatipal and Coumatipal. As there may be some drug interactions. Report any allergic reactions – skin rash, hives, swelling of the face, lips, tongue or throat.
Erythromycin Ophthalmic Ointment	Antibacterial	prophylaxis for infection	apply 1cm V ribbon in eyes x1 w/in 24h after birth	<input checked="" type="radio"/> Y N	n/a	n/a	<ol style="list-style-type: none"> Common reactions include ocular irritation, erythema, and stinging. Contraindicated if there is a hypersensitivity to this med. Used to prevent infections that can cause blindness. Call HCP if you notice allergy's reaction to med.
Engerix B	Vaccine	Hep B vaccine	newborn: 0.5ml IM x1 @ birth	<input checked="" type="radio"/> Y N	n/a	anaphylaxis Stevens Johnson syndrome thrombocytopenia neurotoxicity seizures meningitis Encephalo-bone syndrome osteopenia	<ol style="list-style-type: none"> Common side effects are fatigue, fever, dizziness and headache. educate pt on vaccine, give vaccine packet and get consent signed before admin. Contraindicated if pt has hypersensitivity to yeast. Must adhere to dosing schedule for newborn!
Hepatitis B Immune Globulin		Prevent consequent Chronic liver and Carcinoma of the liver	0.5ml IM w/in 12h of birth	<input checked="" type="radio"/> Y N	n/a	back pain headache muscle aches nausea pain @ injection site	<ol style="list-style-type: none"> educate pt on injection and get consent inform pt that pain @ injection site is common report any nausea or vomiting Newborns may experience soreness, swelling or redness where vaccine was given.
				Y N			<ol style="list-style-type: none">