

Student Name: Joanna Laguna

Unit: Ped/Floor

Pt. Initials: AP

Date: 08/30/23

Pediatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: Povidone-iodine

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Therapeutic Range?	IVP – List solution to dilute and rate to push. IVPB – concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				Is med in therapeutic range? If not, why?			
levetiracetam	anticonvulsant	tx. seizures and help control it	180mg IV Q 12 H	5-55 mg/kg/day Yes, in range	0.9% NaCl (10mg/mL) 72 mL/hr over 15 mins.	Angioedema Hallucinations Steven-Johnson syndrome Toxic epidermal necrolysis	1. Avoid tasks requiring alertness. 2. Report if seizures are different or worsen. 3. Do not suddenly stop use. Slowly taper off. 4. Know that risks are higher for children.
famotidine	H2 blocker/antagonist	prophylaxis for peptic ulcer	5.6 mg 6 Tube BID	0.5-1 mg/kg/day max: 40 mg/day Yes, in range		Fat, abnormal HR Dizziness or passing out Allergy rxn: wheezing tight chest, swell	1. PO meds might be prevented to be absorbed. Take in another time. 2. Do not give for longer than prescribed. 3. Take as directed and complete course. 4. May cause drowsiness or dizziness.
levothyroxine	Thyroid preparations	Thyroid supplementation in hypothyroidism	37.5 mcg 6 Tube daily	5-6 mcg/kg/day No, however children with chronic hypothyroid start at 25 mcg/day		Weight loss Hyperthyroidism	1. Follow-ups are important to assess effectiveness. 2. Take at least 4 hrs apart from antacids. 3. Report S/S of overdose: chest pain, tremors, and other hyperthyroidism manifestation. 4. Temporary, partial hair loss may occur.
							1. 2. 3. 4.
							1. 2. 3. 4.