Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day
Student Name: Christopher Tsai **Date: 9/19/2023**
Patient Allergies: KNA
Primary IV fluid and rate: N/A

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic): N/A

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
Cinacalcet (Sensipar)	30 mg PO Daily		- Cinacalcet decreases levels of parathyroid hormone (PTH), calcium, and phosphorous in the body	1. You should not take cinacalcet if you have low levels of calcium in your blood 2. Cinacalcet works best if you take it with food or shortly after eating a meal 3. your blood will need to be tested 1 week after you start taking cinacalcet or whenever your dose is changed. You may also need blood tests on a regular basis during treatment.
Gabapentin (Neuontin)	300 mg PO TID		- used to treat partial seizures, nerve pain from shingles and restless leg syndrome.	1. Avoid driving or hazardous activity until you know how gabapentin will affect you. 2. Some people have thoughts about suicide while taking seizure medicine. 3. You should not stop taking gabapentin suddenly. Stopping suddenly may cause increased seizures.
Levetiracetam (Keppra)	300 mg PO BID		- used to treat certain types of seizures in people with epilepsy, including partial onset seizures, myoclonic seizures, and tonic-clonic seizures	1. Some people have thoughts about suicide while taking levetiracetam. 2. Levetiracetam oral should take by mouth at the same time each day. 3. Your kidney function need to be

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day
Student Name: Christopher Tsai **Date: 9/19/2023**
Patient Allergies: KNA
Primary IV fluid and rate: N/A

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic): N/A

				tested.
Nifedipine (Adalet)	90 mg PO Daily		- calcium channel blockers. It works by relaxing the muscles of your heart and blood vessels and reduce the BP.	1. Do not use if you have had a heart attack within the past 2 weeks 2. Stopping suddenly may make your condition worse. 3. Your blood pressure will need to be checked often and you may need other blood tests at your doctor's office.
Sevelamer Carbonate (Renevela)	240 mg PO TID		- Sevelamer is used to control phosphorus levels in people with chronic kidney disease who are on dialysis	1. You should not take sevelamer if you have a bowel obstruction 2. Take sevelamer with meals. 3. While using sevelamer, you may need frequent blood tests.
Atorvastatin (Lipitor)	40 mg PO Daily		- It works by slowing the production of cholesterol in the body to decrease the amount of cholesterol that may build up on the walls of the arteries and block blood flow to the heart, brain	1. Atorvastatin is used together with diet, weight loss, and exercise to reduce the risk of heart attack and stroke 2. Atorvastatin can cause the breakdown of muscle tissue, which can lead to kidney failure 3. Avoid eating foods that are high in fat or cholesterol. Atorvastatin will not be as effective in lowering your cholesterol if you do not follow a cholesterol-lowering diet plan.

Patient Name: Christopher Warren

Date/Time: 12/12/2023 Shift for IV fluid/ink (hospital type, isotonic, hypotonic, hypertonic) N/A

Patient Allergies: KNA

Primary IV fluid and rate: N/A

Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Medication reference: https://www.drugs.com/