

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
ceftazidime	1g IVPB Every 8 hours		Antibiotic use to treat bacterial infection	1.call your doctor if you feel bloating. 2.report any change in skin color 3.call your doctor if severe, watery, or bloody diarrhea occur. 4.report abdominal or stomach cramps or tenderness.
Regular insulin	Subcutaneous Sliding scale 131-200 give 2 units. 201-250 give 4 units. 251-300 give 6 units.		Used to lower level of glucose in the blood. Hep improve blood sugar control in adult.	1.Report if fast heartbeat occurs. 2.call your doctor if experienced blurred vision. 3.report difficult, burning, or painful urination. 4.call your doctor if you experience shakiness.

Allergies: NKA

Primary IV fluid and rate: 0.9% NS 500 mL bolus

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Medication reference: Drug.com

Nursing diagnosis and nursing interventions

- 1- High respiratory rate noted: check the O2 sat, instruct the patient to take a slow deep breath in through the nose out through the mouth.
- 2- Diabetic with an open wound foot ulcer: assess the patient's wound, administer antibiotic as prescribe.