

IM 7 Math Module

Complete the required math problems and submit to Math drop box
 Name Tiquisha Mitchell

1. Infuse 1 gram of a medication over 45 minutes. The drug is supplied as 1gram/50ml.

The drip factor is 15. How many gtt/min will you infuse?

$$\frac{50\text{ml} \times 15\text{gtt/ml}}{45\text{min}} = \frac{750\text{gtt}}{45\text{min}} = 16.6 = \boxed{17\text{gtt/min}}$$

2. The physician writes an order to give 1000mL of intravenous fluid over 8hrs. How many mL/hr will you infuse?

$$\frac{1000\text{ml}}{8\text{hr}} = \boxed{125\text{ml/hr}}$$

3. Infuse 1.5 gram of a medication over 3 hours. The drug is supplied as 1.5 gram/250mL.

$$3\text{hr} \times 60\text{min} = 180\text{min}$$

The drip factor is 15. How many gtt/min will you infuse?

$$\frac{250\text{ml} \times 15\text{gtt/ml}}{180\text{min}} = \frac{3750\text{gtt}}{180\text{min}} = 20.83 = \boxed{21\text{gtt/min}}$$

4. An order has been written to give 1 gram of a medication over 30 minutes. The drug is supplied as 1 gram/50mL. The gtt factor is 60. How many gtt/min will you infuse?

$$\frac{50\text{ml} \times 60\text{gtt/ml}}{30\text{min}} = \frac{3000\text{gtt}}{30\text{min}} = \boxed{100\text{gtt/min}}$$

5. The nurse is to give 500mg IV of a medication over 1 hr. The drug is supplied as 1gram/250mL. The gtt factor is 15. How many gtt/min will you infuse?

$$\frac{250\text{ml} \times 15\text{gtt/ml}}{60\text{min}} = \frac{3750\text{gtt}}{60\text{min}} = 62.5 = \boxed{63\text{gtt/min}}$$

6. An order is received for 75mcg IV of a medication now. The drug is supplied as 100mcg/2mL. How many mL will you give?

$$\frac{D}{H} \times V = \frac{75\text{mcg}}{100\text{mcg}} \times 2\text{ml} = 0.75\text{mcg} \times 2\text{ml} = \boxed{1.5\text{ml}}$$

7. Infuse 1000 mLs of intravenous fluid over 4 hrs. How many mL/hr will you set on the pump?

$$\frac{1000\text{ml}}{4\text{hrs}} = \boxed{250\text{ml/hr}}$$

IM 7 Math Module

Complete the required math problems and submit to Math drop box

8. The patient is to receive 5mg of a medication. The drug is supplied as 20mg/5mL. How many mL will you give? (Do not round your final answer)

$$\frac{D}{H} \times V = \frac{5\text{mg}}{20\text{mg}} \times 5\text{mL} = 0.25\text{mg} \times 5\text{mL} = \boxed{1.25\text{mL}}$$

9. The order is to give 2mg IV of a medication now. The drug is supplied as 10mg/mL. How many mL will you give?

$$\frac{D}{H} \times V = \frac{2\text{mg}}{10\text{mg}} \times 1\text{mL} = 0.2\text{mg} \times 1\text{mL} = \boxed{0.2\text{mL}}$$

10. Infuse 500mg IV of a medication over 8 hrs. The drug is supplied as 500mg/100mL. The drip factor is 60. How many gtt/min will you infuse?

$$\frac{100\text{mg} \times 60\text{gtt/hr}}{5\text{hr} \times 60\text{min}} = \frac{1,200\text{gtt}}{480\text{min}} = 2.5 = \boxed{13\text{gtt/min}}$$

11. The patient is receiving an intravenous medication currently infusing at 142 mL/hr. The IV bag of reads 50 mg in 500 mL D5W. The patient weighs 175 lbs. How many mcg/kg/min are infusing? Round to the nearest tenth.

$$\frac{50\text{mg}}{500\text{mL}} \Rightarrow 0.1\text{mg/mL} \text{ or } 100\text{mcg/mL} \quad \frac{175\text{lb}}{2.2\text{kg}} = 79.5\text{kg} \quad \frac{142\text{mL}}{60\text{min}} = 2.36\text{mL/min} \times 100\text{mcg/mL} = 236\text{mcg} \div 79.5\text{kg} = 2.97$$

12. The physician has ordered a medication that states to start at 1 mcg/kg/min, and titrate as needed. The IV bag of medication contains 250 mg in 500 mL D5W. The patient weighs 70 kg. How many mL/hr should the IV pump be set at to achieve the starting dose? Round to the nearest whole number.

$$\frac{1\text{mcg}}{\text{kg} \times \text{min}} \times 70\text{kg} = 70\text{mcg/min} \times 60\text{min} = 4,200\text{mcg/hr} \quad \frac{250,000\text{mcg}}{500\text{mL}} = 500\text{mcg/mL} \quad \frac{4,200\text{mcg/hr}}{500\text{mcg/mL}} = 8.4 = \boxed{8\text{mL/hr}}$$

13. The patient is currently receiving a medication at 12 mL/hr. The bottle reads 100 mg in 250 mL D5W. How many mcg/min is the patient receiving?

$$\frac{100\text{mg}}{250\text{mL}} = 0.4\text{mg/mL} \times 12\text{mL/hr} = 4.8\text{mg/hr} = \frac{4,800\text{mcg/hr}}{60\text{min}} = \boxed{80\text{mcg/min}}$$

14. The physician orders a heparin infusion at 500 units/hr. The IV bag of medication reads 25,000 units in 250 mL D5W. How many mL/hr should be showing on the IV pump?

$$\frac{D}{H} \times V = \frac{500\text{units/hr}}{25,000\text{units}} \times 250\text{mL} = 0.02\text{units/hr} \times 250\text{mL} = \boxed{5\text{mL/hr}}$$

IM 7 Math Module

Complete the required math problems and submit to Math drop box

15. The physician has ordered a medication to start at 2 mcg/kg/min. The patient weighs 165 lbs. The IV bag reads 800 mg in 500 mL D5W. What rate would the nurse set on the infusion pump? Round to the nearest tenth.

Handwritten work for problem 15:

$$2 \text{ mcg/kg/min} \times 165 \text{ kg} \rightarrow 330 \text{ mcg/min} = 330 \text{ mcg} \times \frac{1000 \text{ mcg}}{1 \text{ mg}} = 330,000 \text{ mcg/hr}$$

$$\frac{800 \text{ mg}}{500 \text{ mL}} = 1.6 \text{ mg/mL} = 1600 \text{ mcg/mL}$$

$$\frac{330,000 \text{ mcg/hr}}{1600 \text{ mcg/mL}} = 206.25 \text{ mL/hr}$$

OR

$$\frac{330 \text{ mcg/min}}{1.6 \text{ mg/mL}} = 206.25 \text{ mL/hr}$$

16. The physician in the previous questions has now written an order to increase the medication to 4 mcg/kg/min. Using the information in the previous question, what rate would the nurse set on the IV pump? Round to the nearest tenth.

Handwritten work for problem 16:

→ doubled. $5.625 \times 2 = 11.25 \rightarrow \boxed{11.3 \text{ mL/hr}}$

↓ nearest whole = 11

17. The patient is on a regular insulin drip infusing at 5 units/hr. The bag is labeled 100 units in 250 mL NS. At what rate should the pump be infusing? Round to the nearest whole number.

Handwritten work for problem 17:

$$\frac{0}{\text{hr}} \times \text{IV} = \frac{5 \text{ units}}{100 \text{ units}} \times 250 \text{ mL} = 12.5 \text{ mL/hr} \rightarrow \boxed{13 \text{ mL/hr}}$$

18. The patient is on a medication drip infusing at 35 mL/hr. The label reads 400 mg in 500 mL D5W. The patient weighs 62 kg. How many mcg/kg/min is the patient receiving? Round to the nearest tenth.

Handwritten work for problem 18:

$$\frac{400 \text{ mg}}{500 \text{ mL}} = 0.8 \text{ mg/mL} = 28 \text{ mg/hr}$$

$$\frac{28 \text{ mg/hr}}{60 \text{ min}} = 0.466 \text{ mg/min}$$

$$\frac{0.466 \text{ mg/min}}{62 \text{ kg}} = 7.5 \text{ mcg/kg/min}$$

↓ nearest tenth = 7.5

19. The physician has ordered 1 gram IV of a medication over 30 minutes. Pharmacy has sent an IV bag labeled 1 gram in 50 mL D5W. The IV tubing delivers 15 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

Handwritten work for problem 19:

$$\frac{50 \text{ mL} \times 15 \text{ gtt/mL}}{30 \text{ min}} = \frac{750 \text{ gtt}}{30 \text{ min}} = \boxed{25 \text{ gtt/min}}$$

20. The patient is to receive 400 mg IV of a medication over 1 hour. You receive an IV bag from the pharmacy labeled 400 mg in 100 mL D5W. The IV tubing delivers 12 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

Handwritten work for problem 20:

$$\frac{100 \text{ mL} \times 12 \text{ gtt/mL}}{60 \text{ min}} = \frac{1200 \text{ gtt}}{60 \text{ min}} = \boxed{20 \text{ gtt/min}}$$