

IM 7 Math Module

Complete the required math problems and submit to Math drop box
Name Sarah Shinsato

1. Infuse 1 gram of a medication over 45 minutes. The drug is supplied as 1 gram/50ml.

The drip factor is 15. How many gtt/min will you infuse? $\frac{50 \text{ mL} \times 15}{45} = 17 \text{ gtt/min}$

2. The physician writes an order to give 1000mL of intravenous fluid over 8hrs. How

many mL/hr will you infuse? $\frac{1000 \text{ mL}}{8 \text{ h}} = 125 \text{ mL/hr}$

3. Infuse 1.5 gram of a medication over 3 hours. The drug is supplied as 1.5 gram/250mL.

The drip factor is 15. How many gtt/min will you infuse? $\frac{250 \text{ mL} \times 15}{180 \text{ m}} = 20.8 \rightarrow 21 \text{ gtt/m}$

4. An order has been written to give 1 gram of a medication over 30 minutes. The drug is supplied as 1 gram/50mL. The gtt factor is 60. How many gtt/min will you infuse?

$$\frac{50 \text{ mL} \times 60}{30 \text{ min}} = \frac{3000}{30} = 100 \text{ gtt/min}$$

5. The nurse is to give 500mg IV of a medication over 1 hr. The drug is supplied as 1gram/250mL. The gtt factor is 15. How many gtt/min will you infuse?

$$\frac{500 / 250 \text{ mL}}{1 \text{ hr}} \rightarrow \frac{125 \text{ mL} \times 15}{60 \text{ m}} = 31.25 \rightarrow 31 \text{ gtt/min}$$

6. An order is received for 75mcg IV of a medication now. The drug is supplied as

100mcg/2mL. How many mL will you give? $\frac{75 \text{ mcg} \times 2 \text{ mL}}{100 \text{ mcg}} = \frac{150}{100} = 1.5 \text{ mL}$

7. Infuse 1000 mLs of intravenous fluid over 4 hrs. How many mL/hr will you set on the pump?

$$\frac{1000 \text{ mL}}{4 \text{ h}} = 250 \text{ mL/h}$$

IM 7 Math Module

Complete the required math problems and submit to Math drop box

8. The patient is to receive 5mg of a medication. The drug is supplied as 20mg/5mL. How many mL will you give? (Do not round your final answer)

$$\frac{5\text{mg} \times 5\text{mL}}{20\text{mg}} = 1.25\text{mL}$$

9. The order is to give 2mg IV of a medication now. The drug is supplied as 10mg/mL. How many mL will you give?

$$\frac{2\text{mg} \times 1}{10\text{mg}} = 0.2\text{mL}$$

10. Infuse 500mg IV of a medication over 8 hrs. The drug is supplied as 500mg/100mL. The drip factor is 60. How many gtt/min will you infuse?

$$\frac{100\text{mL} \times 60}{480} = 12.5 \text{ gtt/min}$$

• x.

$$13 \text{ gtt/min}$$

11. The patient is receiving an intravenous medication currently infusing at 142 mL/hr. The IV bag of reads 50 mg in 500 mL D5W. The patient weighs 175 lbs. How many mcg/kg/min are infusing? Round to the nearest tenth.

$$\frac{100\text{mcg} \times 142\text{mL/h}}{79.5 \times 60} = \frac{14,200}{4,770}$$

$$3 \text{ mcg/kg/min}$$

12. The physician has ordered a medication that states to start at 1 mcg/kg/min, and titrate as needed. The IV bag of medication contains 250 mg in 500 mL D5W. The patient weighs 70 kg. How many mL/hr should the IV pump be set at to achieve the starting dose? Round to the nearest whole number.

$$\frac{70\text{kg} \times 1\text{mcg} \times 60\text{min}}{500} = 8.4 \rightarrow 8\text{mL/h}$$

13. The patient is currently receiving a medication at 12 mL/hr. The bottle reads 100 mg in 250 mL D5W. How many mcg/min is the patient receiving?

$$\frac{1}{400}$$

$$\frac{400\text{mcg/mL} \times 12\text{mL/hr}}{60\text{min/hr}} = 80\text{mcg/min}$$

14. The physician orders a heparin infusion at 500 units/hr. The IV bag of medication reads 25,000 units in 250 mL D5W. How many mL/hr should be showing on the IV pump?

$$\frac{\checkmark}{100\text{u/mL}}$$

$$\frac{500\text{u/hr}}{100\text{u/mL}} = 5\text{mL/h}$$

IM 7 Math Module

Complete the required math problems and submit to Math drop box

- 75 15. The physician has ordered a medication to start at 2 mcg/kg/min. The patient weighs 165 lbs. The IV bag reads 800 mg in 500 mL D5W. What rate would the nurse set on the infusion pump? Round to the nearest tenth.

∴
• x •

150 mcg/min

$$1.6 \text{ mg} = 1,600 \text{ mcg} \quad \frac{75 \text{ kg} \times 2 \text{ mcg} \times 60 \text{ min}}{1,600 \text{ mcg}} = 5.6$$

6 mL/hr

16. The physician in the previous questions has now written an order to increase the medication to 4 mcg/kg/min. Using the information in the previous question, what rate would the nurse set on the IV pump? Round to the nearest tenth.

$$\frac{75 \text{ kg} \times 4 \text{ mcg} \times 60}{1,600 \text{ mcg}} = 11.3 \text{ mL/h}$$

17. The patient is on a regular insulin drip infusing at 5 units/hr. The bag is labeled 100 units in 250 mL NS. At what rate should the pump be infusing? Round to the nearest whole number.

$$\frac{5 \text{ u/h}}{0.4} = 13 \text{ u/hr}$$

18. The patient is on a medication drip infusing at 35 mL/hr. The label reads 400 mg in 500 mL D5W. The patient weighs 62 kg. How many mcg/kg/min is the patient receiving? Round to the nearest tenth.

$$\frac{800 \text{ mcg/mL} \times 35 \text{ mL/hr}}{62 \text{ kg} \times 60 \text{ min}} = \frac{28,000}{3,720} = 7.5 \text{ mcg/kg/hr}$$

19. The physician has ordered 1 gram IV of a medication over 30 minutes. Pharmacy has sent an IV bag labeled 1 gram in 50 mL D5W. The IV tubing delivers 15 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

$$\frac{50 \text{ mL} \times 15 \text{ gtt/mL}}{30 \text{ min}} = 25 \text{ gtt/min}$$

20. The patient is to receive 400 mg IV of a medication over 1 hour. You receive an IV bag from the pharmacy labeled 400 mg in 100 mL D5W. The IV tubing delivers 12 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

$$\frac{100 \text{ mL} \times 12 \text{ gtt/mL}}{60 \text{ min}} = 20 \text{ gtt/min}$$