

## IM 7 Math Module

Complete the required math problems and submit to Math drop box

8. The patient is to receive 5mg of a medication. The drug is supplied as 20mg/5mL. How many mL will you give? (Do not round your final answer)

$$\frac{20 \text{ mg}}{5 \text{ mL}} \quad \frac{5 \text{ mg}}{x \text{ mL}} = 1.25 \text{ mL/hr}$$

9. The order is to give 2mg IV of a medication now. The drug is supplied as 10mg/mL.

How many mL will you give?  $\frac{10 \text{ mg}}{1 \text{ mL}} \quad \frac{2 \text{ mg}}{x \text{ mL}} = 0.2 \text{ mL}$

10. Infuse 500mg IV of a medication over 8 hrs. The drug is supplied as 500mg/100mL. The drip factor is 60. How many gtt/min will you infuse?

$$8 \times 60 \text{ min} = 480 \quad 40 \times 100 \mid 480 = 12.5 \rightarrow 13 \text{ gtt/min}$$

11. The patient is receiving an intravenous medication currently infusing at 142 mL/hr. The IV bag of reads 50 mg in 500 mL D5W. The patient weighs 175 lbs. How many mcg/kg/min are infusing? Round to the nearest tenth.

$$\frac{142 \times 100}{79.5 \times 60}$$

$$175 = 79.5 \text{ kg} \quad 50 \text{ mg} \mid 500 \text{ mL} = 0.1 \text{ concentration} = 100 \text{ mcg}$$

$$142 \text{ mL/hr} \quad 14200 \mid 4770 = 2.9 \text{ mcg/kg/min}$$

12. The physician has ordered a medication that states to start at 3 mcg/kg/min, and titrate as needed. The IV bag of medication contains 250 mg in 500 mL D5W. The patient weighs 70 kg. How many mL/hr should the IV pump be set at to achieve the starting dose? Round to the nearest whole number.

$$500 \text{ mL} \mid 60 \text{ min} = 8.3 \text{ mL/hr}$$

13. The patient is currently receiving a medication at 12 mL/hr. The bottle reads 100 mg in 250 mL D5W. How many mcg/min is the patient receiving?

$$\frac{1000 \text{ mcg}}{250 \text{ mL}} = 400 \text{ mcg} \times 12 = \frac{4800}{60 \text{ min}} = 80 \text{ mcg/min}$$

14. The physician orders a heparin infusion at 500 units/hr. The IV bag of medication reads 25,000 units in 250 mL D5W. How many mL/hr should be showing on the IV pump?

$$\frac{500 \text{ units/hr}}{100 \text{ concentration}} = 5 \text{ mL/hr}$$

## IM 7 Math Module

Complete the required math problems and submit to Math drop box

15. The physician has ordered a medication to start at 2 mcg/kg/min. The patient weighs 165 lbs. The IV bag reads 800 mg in 500 mL D5W. What rate would the nurse set on the infusion pump? Round to the nearest tenth.

$$75 \text{ kg} \quad 1.6 \text{ concentration} \quad \frac{75 \times 2 \times 60 \text{ min}}{1000} = 5.4 \text{ mL/hr}$$

16. The physician in the previous questions has now written an order to increase the medication to 4 mcg/kg/min. Using the information in the previous question, what rate would the nurse set on the IV pump? Round to the nearest tenth.

$$\frac{75 \times 4 \times 60}{1000} = 11.3 \text{ mL/hr}$$

17. The patient is on a regular insulin drip infusing at 5 units/hr. The bag is labeled 100 units in 250 mL NS. At what rate should the pump be infusing? Round to the nearest whole number.

$$\frac{5 \text{ units/hr}}{0.4} = 12.5 = 13 \text{ mL/hr}$$

18. The patient is on a medication drip infusing at 35 mL/hr. The label reads 400 mg in 500 mL D5W. The patient weighs 62 kg. How many mcg/kg/min is the patient receiving? Round to the nearest tenth.

$$62 \text{ kg} \quad 0.8 \times 1000 = 800 \text{ mcg} \quad \frac{800 \text{ mcg}}{100 \text{ min}} \times 62 \text{ kg} = 7.5 \text{ mcg/kg/min}$$

19. The physician has ordered 1 gram IV of a medication over 30 minutes. Pharmacy has sent an IV bag labeled 1 gram in 50 mL D5W. The IV tubing delivers 15 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

$$50 \times 15 \div 30 = 25 \text{ gtt/min}$$

20. The patient is to receive 400 mg IV of a medication over 1 hour. You receive an IV bag from the pharmacy labeled 400 mg in 100 mL D5W. The IV tubing delivers 12 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

$$100 \times 12 \div 60 = 20 \text{ gtt/min}$$