

Student Name: Saul Benavidez

Unit: 59

Pt. Initials: K6

Date: 2/26/24

Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: NKA

| Primary IV Fluid and Infusion Rate (ml/hr) | Circle IVF Type | Rationale for IVF | Lab Values to Assess Related to IVF | Contraindications/Complications |
|--|--|-------------------|-------------------------------------|---------------------------------|
| 0.9% Sodium Chloride | <u>Isotonic</u> / Hypotonic / Hypertonic | Hyponatremia | Cl, Na | Fluid overload |

| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
|---|------------------------------|--------------------|------------------------------------|---|--|---|--|
| Adult multivitamin with minerals and iron | Vitamin | Withhold symptoms | 1 tablet PO daily | <input checked="" type="radio"/> Y N | | Allergic reaction | <ol style="list-style-type: none"> S/S anaphylaxis nausea, abd distention do benefits in nursing stool output |
| Folic acid | Vitamin | Anemia | 1mg oral daily | <input checked="" type="radio"/> Y N | | Allergic sensitization | <ol style="list-style-type: none"> may alter sleep pattern mental depression nausea abdominal distention S/S anaphylaxis |
| Thiamine | Vitamin | withhold symptoms | 100mg oral daily | <input checked="" type="radio"/> Y N | | Alcohol intolerance hypersensitivity | <ol style="list-style-type: none"> S/S anaphylaxis resistance of alcohol on mucous membrane S/S hypotension may cause nausea, abd pain |
| Lorazepam | Sedating | withhold symptoms | 2mg sublingual for this indication | <input checked="" type="radio"/> Y N | NS, D5W, 100mL Rate of injection not to exceed 2mg/min | Respiratory Depression | <ol style="list-style-type: none"> avoid alcohol or other sedatives highly addicting do driving after on medication Go to ER if needed |
| Ondansetron | Antiemetic | N/V | 4mg sublingual | <input checked="" type="radio"/> Y N | NS or D5W Rate of injection over 2 to 5 minutes (OT inhibition) | Affect the heart rhythm (QT prolongation) | <ol style="list-style-type: none"> S/S anaphylaxis may cause dizziness S/S hypotension bradycardia/ECG in respiratory |