

Student Name: Ty Skiles Unit: PICU Pt. Initials: PB Date: 4/13/24

Pediatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: NKPA

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
<u>NS 2 mL/Hr</u>	<u>Isotonic/Hypotonic/Hypertonic</u>	<u>Maintains Proper Fluid Balance</u>	<u>Na⁺ & K⁺</u>	<u>Fluid Overload</u>

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Therapeutic Range?		IVP – List diluent solution, volume, and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				Is med in therapeutic range?	If not, why?			
Albumin	Colloid	Helps pull fluid from interstitial tissue back into the blood stream.	25g IVPB Once	Yes		25g / 500ml 25 mL/Hr.	<ul style="list-style-type: none"> Circulatory collapse Hyperkalemia Hypotension HTN 	<ol style="list-style-type: none"> This medication should not turn the skin pale. There should be no edema or around the IVPB. Pt. should not continue to become more swollen. Monitor creatinine. This medication is nephrotoxic.
Neuraxium	Blocker	This medication is used to paralyze the Pt. by relaxing the skeletal muscle.	4mg/kg IVPB Continuous	Yes		0.4 mg/mL 8.34 mL/Hr.	<ul style="list-style-type: none"> Respiratory Distress Bradycardia Flushing 	<ol style="list-style-type: none"> Pt. should not be able to make muscle movement. This medication should not affect the patient's neuromuscular function later in life. One side of the body should not be more affected than the other. Monitor creatinine. This drug is nephrotoxic.
Clinamycin	Antibiotic	This medication is given to combat bacterial infection.	144mg IVPB TID	Yes		12 mg/mL 24 mL/Hr.	<ul style="list-style-type: none"> C. Diff Nephrotoxicity Thrombocytopenia 	<ol style="list-style-type: none"> Report if Pt. stool has an abnormal odor. Report any bloody stool. WBCs should not decrease to Abnormally low Value. This medication may cause anuria.
Furosemide	Loop Diuretic	This medication increases diuresis & treats fluid overload.	0.5mg/kg IVPB Continuous	Yes		1.39 mg/mL 0.7 mL/Hr.	<ul style="list-style-type: none"> Dehydration Hypovolemia HTN Hypernatremia 	<ol style="list-style-type: none"> Report any seizure like activity. Keep Pt. on EKG to monitor for dysrhythmia. WBCs Neutrophils should not become low. Report facility breath odor.
Flaxium Chloride	Electrolyte	This medication is given to correct abnormally low potassium levels.	6.96mg IVPB PRN	Yes		0.2 m Eq/mL 34.8 mL/Hr.	<ul style="list-style-type: none"> Hypernatremia Hypocalcemia Hypomagnesemia Muscle Weakness 	<ol style="list-style-type: none"> Report any edema that looks like coffee beans. Stomach bloating is abnormal. Muscles should not begin to twitch. Report any seizure like activity.