

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)		Circle IVF Type		Rationale for IVF		Lab Values to Assess Related to IVF		Contraindications/Complications	
Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>									
Click here to enter text.		Click here to enter text.		Click here to enter text.		Click here to enter text.		Click here to enter text.	
Allergies: Click here to enter text.									
Adverse Effects Click here to enter text.									
Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)									
Date: Click here to enter a date.									
Correct Dose? If not, what is correct dose?									
Dose, Route & Schedule									
Therapeutic Reason									
Pharmacologic Classification									
Generic Name									
Unit: Click here to enter text.									
Patient Initials: Click here to enter text.									
Student Name: Brianna Amaya									
Hydroxyzine pamoate	Antihistamine	Anxiety	25mg PO PRN q6	Yes	NA	Common SE: HA, dry mouth, skin rash, drowsiness	1. can cause birth defects in women 2. call dr if you have CP with HA 3. caution when driving can make you sleepy 4. alcohol can increase risk of SE		
lorazepam	Benzodiazepine	Alcohol withdrawal / anxiety	1-2 mg PO PRN q4 hours	Yes	NA	Common SE: sedation, dizzy, drowsiness, weakness, unsteady,	1. do not drink use opioids or other drugs that can slow breathing 2. take as prescribed can become addictive 3. women should not take if pregnant 4. keep out of reach of children		
haldol	antipsychotic	for psychotic symptoms/agitation	10 mg IM PRN q8	Yes	NA	Common SE: HA drowsiness, insomnia, restlessness, dizziness	1. may cause uncontrolled muscle movements (lip smacking, tongue movement, eye movement) 2. long term use can cause irreversible SE 3. never take more than prescribed can cause heart rhythm disorder 4. may take several weeks to take effect		

Orrianna Amaya

Current Medications & PRN for Last 24 Hours

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	Date: Click here to enter a date.	Adverse Effects	Allergies: Click here to enter text.	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
diphenhydramine	antihistamine	management of cholinergic effects of antipsychotic	50 mg IM q8 PRN psychotic symptoms	Yes	NA	SE: constipation, drowsiness, dry eyes, dizzy		1. caution while driving (drowsiness) 2. Children under 2 YOA should not take 3. call dr if tightness in neck or jaw occur 4. take fiber if constipation occur
acetaminophen	Analgesic/antipyretic	Mild-moderate pain	500 mg PO q6 for pain of 1-4 on pain scale	Yes	NA	Common SE: N/V/D, constipation, abdominal pain, HA		1. ETOH may increase risk for liver damage 2. if yellowing of skin or eyes notify HCP (liver damage) 3. do not exceed >4000mg in 24 hrs 4. stomach cramps or pain call HCP (overdose symptom)
olanzapine	antipsychotic	Agitation/psychotic behavior	5 mg PO q6 for agitation	Yes	NA	Common SE: increased appetite, wt gain, HA, dizziness, feeling tired or restless, tremors/shaking constipation		1. Serious drug reaction can affect many parts of the body 2. monitor BGL can cause high blood sugar 3. changes in personality is common with this medication 4. take medication exactly as directed

Adopted: August 2016

+

Student Name: Brianna Amaya
Communication Worksheet – Current Medications & PRN for Last 24 Hours

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	Date:	Allergies:	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
ibuprofen	NSAID	Mild-moderate pain	400 mg PO q6 for pain of 1-4 on pain scale	Yes	NA IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Common SE: N/V, gas, bleeding, dizziness, HA	1. ask HCP if okay to take if on antidepressants 2. tell dr if you take aspirin to prevent MI or CVA 3. don't exceed 800mg in a dose or 3200mg per day 4. take the lowest dose needed to treat condition
risperdal	antipsychotic	Psychotic behavior/bipolar mania	1 mg tablet PO BID	Yes	NA	Common SE: depressed mood, HA, Dizzy, tremors/twitching, agitation, anxiety, restless, dry mouth, wt gain	1. do not use in adults with dementia related psychosis 2. avoid getting too cold or too hot (dehydrated) 3. stop using and notify dr if fever, stiffness, and confusion occur (adverse SE) 4. may temporarily affect fertility in women
Nicotine patch	Stimulant	Nicotine replacement	21 mg transdermal patch daily	Yes	NA	SE: sleeping disturbances, skin redness, itching burning, HA	1. rotate patch site every day 2. seek treatment if allergic reaction occurs 3. if HA occurs try asking for a lower dose patch

Adopted: August 2016

Student Name: Brianna Amaya		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	Allergies: Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
ziprasidone	antipsychoti c	Psychotic behavior/bipol ar disorder/mania	40 mg PO BID	Yes	NA	SE: dizziness, drowsiness, HA, n/v, restlessness, wt gain, tremors	1. get emergency help if signs of allergic reaction or severe skin reaction occur 2. can cause high blood sugar 3. stiff neck may occur 4. tell the HCP other medication you take like antidepressants (contraindicated)
prazosin	Alpha- blocker	Nightmares	1 mg PO QHS	Yes	NA	SE: dizzy, HA, drowsiness, feeling weak or tired, nausea	1. males call HCP if erection lasting longer than 4 hours occurs (damaging) 2. medication can cause low BP 3. do not stop can cause rebound HTN 4. in females can cause diabetes and eclampsia during pregnancy
Nicotine inhaler	Stimulant	Nicotine replacement	10 mg cartridge for inhalation Q2	Yes	NA	SE: dizziness, dry mouth, hiccups, burping, upset stomach, HA, change in taste	1. call dr if pounding or fast heartbeats or fluttering 2. stop using if wheezing tightness in chest or SOB 3. keep our of reach of children can be

Adopted: August 2016

Date: 4-18-14
 Simulation Topic: ETOI with A/Causal
 Observer: Sgt. Amaya
 Primary Nurse: Andrea
 Secondary Nurse: Vicki
 Patient/Family member/s: Valeria

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: Brianna Amaya		Unit:	Patient Initials:	Date:	Allergies:	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IV/PB – List diluent solution, volume, and rate of administration	IV/PB – List concentration and rate of administration	fatal
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter a date.	Click here to enter text.	Click here to enter text.	4. women do nto use if pregnant
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.