

Child & Adolescent Disorders Activity

Autism Spectrum Disorder (ASD)

Attention Deficit Disorder (ADHD)

Oppositional Defiant Disorder (ODD)

Conduct Disorder (CD)

Childhood & Adolescent Depression (Dep)

Which disorder does the symptom, medication or therapy pair with?

May be temporary & precipitated by a traumatic event
Vandalism
Difficult to assess for pain
Psychostimulants
Do not see themselves as defiant
Early Childhood Intervention
Substance use
Interrupts others
Decrease in school performance
Limited facial expression
Breaks the law
Gluten-free/casein free diet
May need verbal & written instructions
Selective serotonin reuptake inhibitors (SSRI's)
Explosive angry outbursts
"Stimming"
Weight loss or weight gain
Echolalia
At greater risk for injury
Disinterest in previously enjoyed activities
Anger management classes
Difficulty completing tasks
Temper tantrums
Seeks revenge
Antipsychotic medications
Demonstrates a lack of remorse
Hopelessness
Absent or delayed speech
See their behavior as a response to unreasonable demands
May have difficulty following instructions
Hippotherapy
Desire to be alone
Talks excessively
May grunt or hum instead of talk
Melatonin for sleep
Bullying
Mood stabilizers
Blames others for disagreements
Applied Behavior Therapy (ABA)
Difficulty sitting still
M-CHAT