

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
D5NS + 20K @ 5 mL/hr	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input checked="" type="checkbox"/>	Fluid and electrolyte replacement	Click here to enter text.	None

Student Name: Jordan Magee		Unit: PICU and Pedi		Patient Initials:		Date: 8/29/2024		Allergies: NKDA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)		
Enoxaparin	Anticoagulant	Prevent blood clots	40mg, Subq, BID	Yes	N/A	Confusion, hemorrhage, anemia, hyperkalemia, anaphylaxis	1. Rotate injection sites to avoid lipodystrophy 2. Notify provider of signs of bleeding or thrombocytopenia 3. Use soft-bristle toothbrush and electric razor to avoid bleeding gums and cutting yourself - due to bleeding risk 4. Avoid OTC products containing aspirin or NSAIDs unless approved by provider		
Famotidine	Antilulcer agent	Prevent stomach ulcers and upset	20mg, IVP, BID	Yes Choose an item.	IVP - Pull up 2 mL, push over 2 minutes	Seizures, insomnia, tinnitus, anorexia, pneumonia	1. Can cause dysrhythmias or QT prolongation so notify physician immediately if any cardiovascular changes/symptoms occur 2. Avoid irritating foods, alcohol, aspirin, NSAIDs, and extreme temperature foods that may irritate the GI system 3. Product must be continued for prescribed time to be effective; do not double dose or take for extended periods		

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jordan Magee		Unit: PICU and Pedi	Patient Initials:		Date: 8/29/2024	Allergies: NKDA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							of time 4. Increase in bulk and fluids in diet to prevent constipation
Acetaminophen	Nonopioid analgesic	Mild to moderate pain	500mg, IVPB, q6hrs	Yes Choose an item.	IVPB - 10mg/mL, infuse over 15 mins	Dyspnea, atelectasis, GI bleeding, renal failure, thrombocytopenia	1. BBW: High doses can cause hepatotoxicity - Max dose is 4,000mg/day; Be aware of other meds than may contain acetaminophen 2. Do not drink alcohol, use herbals, or take OTC products with this drug without approval of prescriber - combinations could interact with the drug, cause stomach irritation, cause liver damage, or cause other unwanted side effects 3. Notify provider if rash or urticaria occur - could indicate hypersensitivity or allergic reactions and product should be stopped 4. Recognize signs of chronic overdose - bleeding, bruising, malaise, fever, sore throat
Baclofen	Skeletal muscle relaxant	Infantile spasms	2.5mg, PO, TID	Yes Choose an item.	N/A	Insomnia, peresthesias, seizures, hypotension,	1. Do not discontinue med quickly - hallucinations, spasticity, or tachycardia can occur; taper off slowly 2. Rise slowly to prevent orthostatic

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jordan Magee		Unit: PICU and Pedi	Patient Initials:		Date: 8/29/2024	Allergies: NKDA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
						tinnitus, hematuria, dyspnea	hypotension 3. Do not take with alcohol or opioids - could cause respiratory depression, suppressed breathing, and dizziness 4. Notify provider of any rash, fever, or respiratory distress - could indicate allergic reaction
Gabapentin	Anticonvulsant	Seizures	100mg, per G-tube, TID	Yes Choose an item.	N/A	Somnolence, aggression, blurred vision, UTI, leukopenia, gout	1. Do not take within 2 hrs of antacid due to risk of decreased absorption 2. Notify provider of suicidal thoughts or feelings, increased depression, panic attacks, hostility, or confusion 3. Do not discontinue abruptly - could cause seizures 4. Do not use alcohol, sedatives, antihistamines, or other products that cause dizziness/sleepiness with this drug
Clobazam	Anticoagulant	Seizures associated with Lennox-Gastaut syndrome	7.5mg, per G-tube, BID	Yes Choose an item.	N/A	Hyperactivity, insomnia, sedation, UTI, URI, pneumonia	1. Do not discontinue product abruptly; taper off slowly 2. Use caution with hazardous activities until response to drug is known 3. Notify provider of changes in behaviors, depression, or changes in mood - can

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jordan Magee		Unit: PICU and Pedi	Patient Initials:		Date: 8/29/2024	Allergies: NKDA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							cause suicidal thoughts  4. Avoid using opioids or any CNS depressants as combination can cause severe respiratory depression, abuse, or dependence
Lorazepam	Sedative, hypnotic	Seizures	1.1mg, IV, every 15 min PRN	Yes Choose an item.	N/A	Confusion, tremors, hallucinations, apnea, cardiac arrest, acidosis	1. Do not discontinue medication abruptly after long-term use - could cause withdrawal; taper off slowly 2. Do not take more than prescribed amount - may be habit forming 3. Rise slowly to minimize orthostatic hypotension and fainting 4. Notify provider immediately of trouble breathing, dizziness, coma, or if no response - could indicate respiratory depression
Lacosamide	Anticonvulsant	Seizures	50mg, per G-tube, BID	Yes Choose an item.	N/A	Syncope, tremors, orthostatic hypotension, palpitations, hepatitis, anemia	1. Do not discontinue product abruptly; taper over 1 week as seizures may occur 2. Notify provider of any rash, fever, fatigue, yellowing of skin, eyes, dark urine - may be hypersensitivity reaction 3. Avoid hazardous activities until stabilized on product - notify provider of

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jordan Magee		Unit: PICU and Pedi	Patient Initials:		Date: 8/29/2024	<b>Allergies:</b> NKDA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							blurred vision, nausea, dizziness, or syncope  4. Carry emergency ID stating product use
Phenobarbital	Barbituate, antiseizure	Seizures	32.4mg, per G-tube, BID	Yes Choose an item.	N/A	Dizziness, double vision, depression, bradycardia, shallow breathing	1. Do not stop medication abruptly - could cause seizures to worsen 2. Use medication exactly as prescribed to prevent/lower the risk of addiction 3. Notify provider of any unusual/sudden changes in your mood, thoughts, or behavior - could cause suicidal behavior 4. Notify provider of any rash, fever, swollen lymph nodes, itching/swelling - could indicate allergic reaction
	Click here to enter text.	Click here to enter text.		Choose an item. Choose an item. Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item. Choose an item. Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text.

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jordan Magee		Unit: PICU and Pedi	Patient Initials:		Date: 8/29/2024	<b>Allergies:</b> NKDA	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				here to enter text.			4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item. Choose an item.Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item. Choose an item.Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item. Choose an item.Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.