

## Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
Click here to enter text.	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>	Click here to enter text.	Click here to enter text.	Click here to enter text.

Student Name: Makenna Pate		Unit: Click here to enter text.	Patient Initials: Click here to enter text.	Date: Click here to enter a date.	Allergies: Click here to enter text.		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Hydroxyzine pamoate	Antihistamine	Anxiety	25mg PO PRN q6	Choose an item. Click here to enter text.	Click here to enter text.	HA and chest pain, tachycardia	<ol style="list-style-type: none"> <li>1. May cause drowsiness or dizziness</li> <li>2. Rinse mouth frequently and sugarless gum to help with dry mouth</li> <li>3. May cause birth defects, so verify pt is not pregnant</li> <li>4. Stop medication if you experience tremors or seizures.</li> </ol>
lorazepam	Benzodiazepine	Alcohol withdrawal / anxiety	1-2 mg PO PRN q4 hours	Choose an item. Click here to enter text.	Click here to enter text.	Severe and sudden drowsiness	<ol style="list-style-type: none"> <li>1. Do not suddenly stop taking without consulting health care professional.</li> <li>2. Report any unusual movements, responses or expressions</li> <li>3. Educate pt about possible suicidal ideations</li> <li>4. Misuse of this drug can cause addiction or overdose.</li> </ol>
haldol	antipsychotic	for psychotic symptoms/agitation	10 mg IM PRN q8	Choose an item. Click here to enter text.	Click here to enter text.	Tachycardia and uncontrolled muscle movements	<ol style="list-style-type: none"> <li>1. May take several weeks to obtain desired effects</li> <li>2. Educate pt to avoid sun exposure to prevent photosensitivity reactions.</li> <li>3. Report chest pain immediately</li> <li>4. Avoid alcohol and other CNS depressants with this medication</li> </ol>
diphenhydramine	antihistamine	management of cholinergic effects of antipsychotic	50 mg IM q8 PRN psychotic symptoms	Choose an item. Click here to enter text.	Click here to enter text.	Constipation, dry mouth, and confusion	<ol style="list-style-type: none"> <li>1. Alcohol may potentiate drowsiness</li> <li>2. Avoid driving or operating machinery until reaction to medication is known</li> <li>3. Educate importance of additional therapies for muscle spasms</li> <li>4. Importance of routine follow ups</li> </ol>
acetaminophen	Analgesic/antipyretic	Mild-moderate pain	500 mg PO q6 for pain of 1-4 on pain scale	Choose an item. Click here to enter text.	Click here to enter text.	N/V, constipation	<ol style="list-style-type: none"> <li>1. Chronic excessive use may lead to hepatotoxicity, renal and cardiac damage</li> <li>2. Consult health care professional if discomfort or fever not relieved by routine dose</li> <li>3. Educate about stomach pain or loss of appetite</li> <li>4. Call provider about redness or spreading rash</li> </ol>
olanzapine	antipsychotic	Agitation/psychotic behavior	5 mg PO q6 for agitation	Choose an item. Click here	Click here to enter text.	Orthostatic hypotension, tachycardia or	<ol style="list-style-type: none"> <li>1. Educate pt about possibility of extrapyramidal symptoms</li> <li>2. Caution pt to avoid driving as med</li> </ol>

				to enter text.		bradycardia, chest pain	alters alertness 3. Notify provider of suicidal thoughts or thoughts of dying 4. Educate on limiting sun exposure
ibuprofen	NSAID	Mild-moderate pain	400 mg PO q6 for pain of 1-4 on pain scale	Choose an item. Click here to enter text.	Click here to enter text.	SOB, changes in vision	1. Caution pt about mixing with alcohol as it increases risk of GI bleeding 2. Contact provider if rash, itching, or vision changes occur 3. Take with food to avoid GI upset 4. Drowsiness or dizziness may occur
risperdal	antipsychotic	Psychotic behavior/bipolar mania	1 mg tablet PO BID	Choose an item. Click here to enter text.	Click here to enter text.	Back pain, HA, confusion	1. Educate on importance of taking medication as directed 2. Notify provider of suicidal thoughts or thoughts of dying 3. Educate pt to avoid sun exposure to prevent photosensitivity reactions. 4. Emphasize the importance of routine follow up exams to monitor side effects
Nicotine patch	Stimulant	Nicotine replacement	21 mg transdermal patch daily	Choose an item. Click here to enter text.	Click here to enter text. <i>takes 8 hrs for first patch to cause effect.</i>	Burning at patch site, tachycardia	1. May cause drowsiness or dizziness 2. Apply patch at the same time each day 3. Do not wear more than one patch at a time 4. Participate in a smoking cessation program while using this product
ziprasidone	antipsychotic	Psychotic behavior/bipolar disorder/mania	40 mg PO BID	Choose an item. Click here to enter text.	Click here to enter text.	Orthostatic hypotension	1. Change positions slowly to minimize orthostatic hypotension 2. Advise pt of need for continued medical follow up for psychotherapy, eye exams, and lab tests 3. Notify provider of regimen prior to surgery 4. Stop taking medication if sudden onset of dizziness or heart flutters
prazosin	Alpha-blocker	Nightmares	1 mg PO QHS	Choose an item. Click here to enter text.	Click here to enter text.	Chest pain, light headedness	1. Advise pt to weight self twice weekly and assess for feet and ankle swelling 2. Check BP prior to med admin as it can drop BP 3. Educate on additional interventions for hypotension 4. Assess for feet and ankle swelling
Nicotine inhaler	Stimulant	Nicotine replacement	10 mg cartridge for inhalation Q2	Choose an item. Click here to enter text.	Click here to enter text.	Chest tightness, tachycardia	1. Do not use along with other nicotine products 2. Educate on proper usage 3. Avoid eating or drinking for 15 minutes before and after using the inhaler 4. Stop smoking before inhaler use
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item. Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to	Click here to	Click here to	Click here	Choose an	Click here to enter	Click here to	1. Click here to enter text.

enter text.	enter text.	enter text.	to enter text.	item. Click here to enter text.	text.	enter text.	2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
-------------	-------------	-------------	----------------	------------------------------------	-------	-------------	--

Adopted: August 2016