

## Child & Adolescent Disorders Activity

Autism Spectrum Disorder (ASD)

Attention Deficit Hyperactivity Disorder (ADHD)

Oppositional Defiant Disorder (ODD)

Conduct Disorder (CD)

Childhood Depression

Suicide

## Which disorder does the symptom, medication or therapy pair with?

May be temporary & precipitated by a traumatic event  
Vandalism  
Difficult to assess for pain  
Psychostimulants  
Do not see themselves as defiant  
Early Childhood Intervention  
Substance use  
Assessment utilizing SLAP  
Interrupts others  
Decrease in school performance  
Limited facial expression  
Breaks the law  
Gluten-free/casein free diet  
May need verbal & written instructions  
Selective serotonin reuptake inhibitors (SSRI's)  
Explosive angry outbursts  
"Stimming"  
Weight loss or weight gain  
Echolalia  
At greater risk for injury  
May tell their friends about their plan  
Disinterest in previously enjoyed activities  
Anger management classes  
Difficulty completing tasks  
Temper tantrums  
Seeks revenge  
Antipsychotic medications  
Demonstrates a lack of remorse  
Hopelessness  
Absent or delayed speech  
See their behavior as a response to unreasonable demands  
May have difficulty following instructions  
Hippotherapy  
Desire to be alone  
Talks excessively  
May grunt or hum instead of talk  
Melatonin for sleep  
Bullying  
Mood stabilizers  
Blames others for disagreements  
Applied Behavior Therapy (ABA)  
Difficulty sitting still  
M-CHAT  
Behavioral Contracts