

Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

| Generic Name | Dosage with route and schedule | IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration | Patient specific therapeutic reasoning | Patient specific teaching with reasoning |
|--------------|---|--|--|--|
| Ondansetron | Oral tablets | IV dose should not exceed 8mg <ul style="list-style-type: none"> • 2mg/ml | To prevent nausea and vomiting. | <ol style="list-style-type: none"> 1. Do not chew, it will dissolve on tongue. 2. Take one at a time if you are taking more than one. 3. Call before getting up it might give you a headache, making you feel lightheaded or dizzy. 4. Notify if you have severe side effects such as: abdominal pain or muscle spasms |
| Metoprolol | Oral tablets | IV- 5mg/5ml 5mg IV q2min ? | Hypertension Acute MI Congestive heart failure angina | <ol style="list-style-type: none"> 1. If you have asthma, it may cause bronchospasm 2. In diabetic it will mask the symptoms of hypoglycemia 3. Take as directed and do not stop taking abruptly. 4. Call before getting up this might cause some dizziness |
| Levofloxacin | Can be oral 250mg tabs or Iv fluid 25mg/mL vial | IV infusion 250-500mg over 60 min. | Antibiotic used to treat infection | <ol style="list-style-type: none"> 1. Reports s/s of tendonitis, may cause tendon rupture. 2. Report any numbness or tingling in arms, hands, legs, or feet 3. Report headache, |

Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

| | | | | |
|------------|-----------------------------|--|--|--|
| | | | | <p>changes in vision, hallucinations.</p> <p>4. Notify immediately drooping eyelids and unsteady gait.</p> |
| Vancomycin | IV 500mg-10g solutions | 15mg/kg IV over 1-2 hrs. begin this 2hr before first incision. | To prevent/ treat infection from surgery. | <ol style="list-style-type: none"> 1. May cause N/V, notify nurse. 2. DO NOT recommend for pregnant people (BLACK BOX WARNING) 3. Rapid infusion may cause symptoms of wheezing, dyspnea, chest pain, notify the nurse 4. May cause redness or tenderness to the site, notify nurse. |
| Prednisone | Oral, concentrated solution | Not sure if this can be given IV?? | To treat inflammation; arthritis, blood disorders, breathing problems. | <ol style="list-style-type: none"> 1. Take this with food or milk to prevent upsetting your stomach. 2. Measure with appropriate dropper provided 3. If taking 1 dose take it before 9AM 4. Do NOT stop abruptly. |
| | | | | |

Allergies:

Primary IV fluid and rate:

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Medication reference:

Medscape.com , CVPV powerpoint, medicines.org

Since this med sheet if for the OR I left the highlighted questions blank.