

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
Click here to enter text.	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>	Click here to enter text.	Click here to enter text.	Click here to enter text.

<b>Student Name:</b> Click here to enter text.	<b>Unit:</b> Click here to enter text.	<b>Patient Initials:</b> Click here to enter text.	<b>Date:</b> Click here to enter a date.	<b>Allergies:</b> Click here to enter text.
---	---	---	---	--

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Hydroxyzine pamoate	Antihistamine	Anxiety	25mg PO PRN q6	Choose an item.  Click here to enter text.	Click here to enter text.	Abnormal heart beats  Confusion  Tremors	<ol style="list-style-type: none"> <li>1. This medication may make you drowsy, do not drive while taking this medication until you know how it affects you.</li> <li>2. This medication can cause increased anxiety, notify your pcp if you have any changes in mood or behavior.</li> <li>3. Do not take grapefruit with this medication, it can affect how the medication works.</li> <li>4. Avoid alcohol with this medication, it can increase drowsiness.</li> </ol>
lorazepam	Benzodiazepine	Alcohol withdrawal / anxiety	1-2 mg PO PRN q4 hours	Choose an item.  Choose an item. Click here to enter text.	Click here to enter text.	Seizures  Jaundice  Syncope	<ol style="list-style-type: none"> <li>1. Do not take this medication with any other sedation medications, this can cause a decrease in respiratory function.</li> <li>2. This medication should be taken only as prescribed; it can cause a dependence and should only be used for a short time.</li> <li>3. Do not take more than is prescribed, depression of the respiratory system can</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text.		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	<b>Allergies:</b> Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							occur. 4. Do not take with alcohol, the combination can also cause respiratory depression.
haldol	antipsychotic	for psychotic symptoms/agitation	10 mg IM PRN q8	Choose an item.  Choose an item. Click here to enter text.	Click here to enter text.	Acute dystonia Parkinsonism Orthostatic hypotension Tardive dyskinesia	1. Do not drive until you are aware how this medication affects your response time, as it can cause a slower reflex. 2. Never stop abruptly, this can be fatal 3. Only take as prescribed, this medication can be highly addictive 4. Avoid alcohol, this can intensify the effects of this medication.
diphenhydramine	antihistamine	management of cholinergic effects of antipsychotic	50 mg IM q8 PRN psychotic symptoms	Choose an item.  Click here to enter text.	Click here to enter text.	Cardiac arrhythmias Heatstroke Seizures	1. This medication can cause dizziness, use precautions before standing up. 2. Notify your PCP if you have signs of decreased urine output. This medication can cause urinary retention. 3. Notify your PCP with any signs of blurred visions. This medication can cause Diplopia 4. Notify your PCP if you have trouble breathing, this medication can cause thickened bronchial secretions

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text.		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	<b>Allergies:</b> Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
acetaminophen	Analgesic/antipyretic	Mild-moderate pain	500 mg PO q6 for pain of 1-4 on pain scale	Choose an item.  Click here to enter text.	Click here to enter text.	Bloody or black tarry stools  Skin rash, itching or hives  Sudden decrease in urine  Unusual bleeding or bruising	<ol style="list-style-type: none"> <li>1. Take this medication at first signs of pain occur to insure they work best. Waiting any will cause this medication to not work as well.</li> <li>2. Do not take this medication for more than 10 days because this can cause liver damage.</li> <li>3. Do not take with alcohol as this can increase the risk for liver damage.</li> <li>4. Get help immediately if you have signs of increased sweating, diarrhea, stomach pain/cramps or loss of appetite. These can be signs of overdose.</li> </ol>
olanzapine	antipsychotic	Agitation/psychotic behavior	5 mg PO q6 for agitation	Choose an item.  Click here to enter text.	Click here to enter text.	Seizures  Very stiff muscles  Difficulty breathing or swallowing	<ol style="list-style-type: none"> <li>1. This medication can cause hyperglycemia, notify your HCP if you have signs of excessive thirst, excessive urination or excessive hunger.</li> <li>2. If you forget a dose, take as soon as you remember unless it is close to the next dose and skip the missed dose.</li> <li>3. This medication can cause constipation, talk to your PCP about laxatives if this occurs. You can also drink plenty of water and exercise regularly to help with this</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text.		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	<b>Allergies:</b> Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							side effect. 4. Dizziness can occur with this medication, always get up slowly when standing.
ibuprofen	NSAID	Mild-moderate pain	400 mg PO q6 for pain of 1-4 on pain scale	Choose an item.  Choose an item. Click here to enter text.	Click here to enter text.	Trouble breathing  Pain in lower back, arm, throat  Sudden severe headache	1. Ibuprofen can increase risk for bleeding, notify your PCP if you have signs of easily bruising or bleeding. 2. Ibuprofen can increase the risk for ulcers or tears in the gut. Notify your PCP if you have signs of black tarry stool, feel faint or vomiting blood. 3. This medication can increase high blood pressure. Notify your PCP if you have and chest pain, severe headache, nose bleeds or buzzing in the ears. 4. This medication can cause severe skin reactions, get help right away if you have any signs of painful skin that is red or purple that looks burnt or peels off, thickening of the skin, or sores that crust over while taking this medication.
risperdal	antipsychotic	Psychotic behavior/bipolar	1 mg tablet PO	Choose an item.	Click here to enter text.	Neuroleptic malignant	1. This medication can cause a low WBC, notify your PCP as soon as you have any

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text.		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	<b>Allergies:</b> Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
		ar mania	BID	Click here to enter text.		syndrome  Tardive dyskinesia	symptoms of feeling ill. Take precautions when going in public settings to avoid illnesses.  2. Do not drink alcohol while taking this medication, it can intensify the effects of this med.  3. This medication can cause dizziness, get up slowly when standing.  4. Talk to your PCP before taking any over the counter medication, these can interact with this medication.
Nicotine patch	Stimulant	Nicotine replacement	21 mg transdermal patch daily	Choose an item.  Choose an item. Click here to enter text.	Click here to enter text.	Mental mood changes  Chest pain  Trouble speeding  Weaknes on one side of the body	1. Do not apply patch to irritated or cut skin, this can increase irritation and lessen the effect of the medication.  2. Do not suddenly stop this medication, it can cause withdraw symptoms such as nervousness, irritability, and headache.  3. Mild symptoms such as stinging, redness or burning may occur at the site, but notify your PCP if this last longer than 4 days. This may be a sign of allergic reaction  4. It is important to finish this medication

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text.		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	<b>Allergies:</b> Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							in order to be sure it works. If symptoms still occur when finished, notify your PCP.
ziprasidone	antipsychotic	Psychotic behavior/bipolar disorder/mania	40 mg PO BID	Choose an item.  Click here to enter text.	Click here to enter text.	Chest pain  Cardiac arrhythmias  Trouble swallowing	<ol style="list-style-type: none"> <li>1. Serious skin reactions can occur with this medication, notify your PCP immediately if you notice red skin lesions, skin rash, sores, blistering, ulcers or peeling of this skin</li> <li>2. This medication can cause hyperglycemia, notify your PCP if you have signs of increased thirst or increased urination.</li> <li>3. This medication may cause you to become overheated easily, take precautions if you exercise or are exposed to high temps. Drink plenty of water and try to stay out of the sun.</li> <li>4. Notify your PCP if you plan to take OTC medications, these can interact with your medication.</li> </ol>
prazosin	Alpha-blocker	Nightmares	1 mg PO QHS	Choose an item.  Choose an item.Click	Click here to enter text.	First dose hypotension  Hallucinations	<ol style="list-style-type: none"> <li>1. This medication can cause heart palpitations, notify your PCP if you notice excessive amount at one time.</li> <li>2. This medication can cause orthostatic</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text.		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	<b>Allergies:</b> Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				here to enter text.		Priapism	hypotension, when standing do so slowly to avoid dizziness that may cause falling. 3. This medication may cause weakness but if it begins to bother you, notify your PCP 4. Notify your PCP if you decide to take any OTC medications, these medications can interact with each other.
Nicotine inhaler	Stimulant	Nicotine replacement	10 mg cartridge for inhalation Q2	Choose an item.  Click here to enter text.	Click here to enter text.	Rapid, pounding, or irregular heartbeat  Wheezing or trouble breathing  Itching, hives, swelling of the face	1. Do not smoke while taking this medication, it can cause side effects. 2. Notify your PCP if you experience any hearing/vision changes, confusion, stomach pain, or cold sweats. This may be a sign of overdose. 3. Only take the dose that is prescribe. Taking more can cause overdose affects. 4. If you miss a dose, take immediately, unless it is close to the next dose, skip the missed dose.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text.

## Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

<b>Student Name:</b> Click here to enter text.		<b>Unit:</b> Click here to enter text.	<b>Patient Initials:</b> Click here to enter text.		<b>Date:</b> Click here to enter a date.	<b>Allergies:</b> Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				Click here to enter text.			4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item.  Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.