

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>			

Student Name: **Abiguel Guerrero** Unit: **TTHSC** Patient Initials: Date: **11/5/24** Allergies:

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/ Contraindications, Etc.)
Hydroxyzine pamoate	Antihistamine	Anxiety	25mg PO PRN q6			Blurred vision Dry mouth constipation Dizziness	Encourage sugar free hard candy or juice to relieve dry mouth Report mental or mood changes, may result in being a danger to self or others Get medical help immediately if seizures, tachycardia, or fainting occur could indicate overdose
lorazepam	Benzodiazapine	Alcohol withdrawal / anxiety	1-2 mg PO PRN q4 hours			Syncope Nausea Vomiting Tremors	Inform that this medication is highly addictive Do not take with alcohol, leads to severe respiratory depression & death Do not stop abrupt, withdraw is lethal
haldol	antipsychotic	for psychotic symptoms/ agitation	10 mg IM PRN q8			Increased body temp Dry mouth Sedation Urinary retention	Inform that this medication increases the chance of death in older adults This medication causes sedation, refrain from operating machinery Report mood/behavioral changes, could become a danger to self or others Report muscle stiffness, uncontrolled movements, & protruding tongue as this is a sign of Tardive Dyskinesia which is chronic and permanent
diphenhydramine	antihistamine	management of cholinergic effects of antipsychotic	50 mg IM q8 PRN psychotic symptoms			Impaired coordination Epigastric discomfort Headache Drowsiness	Report sudden trouble breathing as this indicates overdose and requires immediate intervention Caution when driving as this medication causes drowsiness Report to HCP if you feel tightness in neck/jaw, or uncontrollable movements as this indicates EPS symptoms which may be permanent

acetaminophen	Analgesic/ antipyretic	Mild-moderate pain	500 mg PO q6 for pain of 1-4 on pain scale			Vomiting Nausea Loss of appetite Fatigue	Report hives, rash, itching, or difficulty breathing as this indicates an allergic reaction Report sweating, nausea, & vomiting as this may indicate overdose & requires immediate intervention Report yellowing of the skin or eyes as this indicates that damage has been done to the liver
olanzapine	antipsychotic	Agitation/ psychotic behavior	5 mg PO q6 for agitation			Constipation Dizziness Dry mouth Headache	Inform that this medication causes an increase risk of death in seniors Report unusual uncontrolled movement of the face or body as this indicates EPS Report difficult swallowing or breathing as this indicates an allergic reaction Drowsiness, slurred speech, & coma indicate overdose has occurred & requires intervention
ibuprofen	NSAID	Mild-moderate pain	400 mg PO q6 for pain of 1-4 on pain scale			Blurred vision Bleeding Kidney disease Black stool	Inform that this medication increases the risk for MI or CVA May cause ulcers, bleeding, or holes in the esophagus, stomach, or intestine Report cyanosis of lips, mouth, or nose as this indicates overdose Do not take anticoagulants with this medication due to the risk for bleeding
risperdal	antipsychotic	Psychotic behavior/ bipolar mania	1 mg tablet PO BID			Tremors Drooling Muscle cramps Blurred vision	This medication causes dizziness & drowsiness, do not want to operate heavy machinery Do not stand up too quickly as fainting spells are common Maintenance of temperature regulation may be a problem This medication may cause thoughts of suicide report worsening mood/ depression
Nicotine patch	Stimulant	Nicotine replacement	21 mg transdermal patch daily			Skin irritation Dizziness Nausea/vomiting Sleep issues	Report difficulty breathing, rash, or swelling as this indicates an allergic reaction Report an abnormal heart rhythm as this may be life threatening Inform that patches must be replaced daily & sites alternated Inform not to smoke while using a patch as overdose can occur
ziprasidone	antipsychotic	Psychotic behavior/ bipolar disorder/mania	40 mg PO BID			Abnormal muscle movements Psychomotor agitation Blurred vision SOB	Inform that this medication causes changes in heart rhythm which can cause fainting Report onset of seizures to HCP immediately Report a rash, hives, or any allergic reaction immediately Inform that this medication can cause Tardive dyskinesia which is chronic and permanent uncontrollable muscle movement

prazosin	Alpha-blocker	Nightmares	1 mg PO QHS			HTN palpitations Hypotension Chest pain	Report breathing problems such as wheezing immediately as this is a result of serious allergic reaction and medication needs to be discontinued immediately
Nicotine inhaler	Stimulant	Nicotine replacement	10 mg cartridge for inhalation Q2			Throat irritation SOB blisters of the mouth Ulcers	Inform to get up slowly due to sudden drop in blood pressure Do not take with alcohol as this medication causes dizziness and drowsiness it could lead to respiratory depression Report heart palpitations and trouble breathing as this may indicate an overdose Inform that this medication may cause sinus pressure and pain May cause tingling/burning in the hands or feet drooling, confusion, and seizures are indicative of an overdose, required immediate intervention