Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Allergies: NKDA

Primary IV fluid and rate: 9% NS KVO. (ISOTONIC)

SOURCES: drugs.com; Davis Drug Guide; Zoloft.com; Mayo Clinic; Cleveland Clinic

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
OFIRMEV	1000MG IVPB	1000 MG IN 100 ML OVER 15 MINUTES. PRN.	GIVEN TO PT FOR PAIN RELATED TO EPIGASTRIC ISSUES. PT HAS MED RX OF MULTIPLE ABDOMINAL SURGERIES. PAIN RATED AS 8/10	PT INFORMED OF THE FOLLOWING: DRUG CAN CAUSE NAUSEA AND VOMITING. ZOFRAN IS ORDERED PRN. DRUG CAN CAUSE CONSTIPATION. PT'S LAST BM WAS ONE DAY PRIOR TO ADMISSION. PT INFORMED THAT SWELLING AND DIFFICULTY BREATHING CAN OCCUR AND TO CALL IMMEDIATELY IF ITCHING OR SWELLING BECOMES NOTICEABLE.
LOVENOX	40 MG SUB Q	N/A	GIVEN TO PT TO PREVENT CLOTS WHILE PATIENT IS NOT MOBILE.	PT INFORMED OF THE FOLLOWING: DO NOT RUB OR SCRATCH THE INJECTION SITE. REPORT ANY UNUSUAL BLEEDING INCLUDING DARK STOOLS OR BRUISING AND BLEEDING GUMS. DO NOT TAKE WITH ASPIRIN OR NEPROXEN.
MORPHINE	4 MG IVP	4 MG IN 1 ML IVP GIVEN OVER 5 MINUTES.	PRESCRIBED FOR PAIN DUE TO EPIGASTRIC ISSUES.	PT INFORMED OF THE FOLLOWING: CAN CAUSE DIFFICULTY BREATHING. CALL IMMEDIATELY IF THIS OCCURS. CAN CAUSE HYPOTENSION, CALL FOR HELP GETTING OUT OF BED OR STANDING. CAN CAUSE CONSTIPATION.

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Allergies: NKDA

Primary IV fluid and rate: 9% NS KVO. (ISOTONIC)

SOURCES: drugs.com; Davis Drug Guide; Zoloft.com; Mayo Clinic; Cleveland Clinic

ZOFRAN	4 MG IV	4 MG IVP GIVEN OVER 2-4 MINS	PRESCRIBED FOR NAUSEA DUE TO PAIN AND EPIGASTRIC ISSUES.	PT INFORMED OF THE FOLLOWING: DIFFICULTY BREATHING OR SWALLOWING CAN OCCUR. CALL IMMEDIATELY. DIZZINESS OR LIGHTHEADEDNESS IS COMMON. CALL FOR HELP BEFORE STANDING OR GETTING OUT OF BED. CAN CAUSE CONSTIPATION.
PROTONIX	40 MG	40 MG IN 10 ML STERILE WATER. GIVEN IVP OVER 2 MINS.	PRESCRIBED FOR ACID REFLUX (PROTON PUMP INHIBITOR)	PT INFORMED OF THE FOLLOWING: AVOID SMOKING, ALCOHOL, AND CARBONATED DRINKS. AVOID HIGH ACID FOODS. SIDE EFFECTS INCLUDE NVD AND HEADACHE. CAN CAUSE DIZZINESS. CALL FOR HELP BEFORE GETTING OUT OF BED.
ZOSYN	3.375 G	3.375 G IN 50ML NS GIVEN IVPB OVER 4 HOURS.	PRESCRIBED FOR INFECTION RELATED TO EPIGASTRIC PAIN.	PT INFORMED OF THE FOLLOWING: DRUG CAN CAUSE NVD, CONSTIPATION, AND STOMACH PAIN. INFORM HCP IMMEDIATELY IF STOMACH PAIN WORSENS. ITCHING, SWELLING, AND HIVES CAN ALSO OCCUR.

Medication reference: Drugs.com, Davis Drug Guide, Mayo Clinic