

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Allergies: NKDA

Primary IV fluid and rate: 9% NS KVO. (ISOTONIC)

SOURCES: drugs.com; Davis Drug Guide; Zoloft.com; Mayo Clinic; Cleveland Clinic

Casodex BICALUTAMIDE	50 MG TABLET	N/A	STOPS TESTOSTERONE FROM ACTING ON CANCER CELLS AND CAUSING THEM TO MULTIPLY.	Casodex can cause birth defects if a child is conceived while the father is on the drug. Continue using birth control for at least 130 days following the last dose. Casodex can also cause you to sunburn more easily. Use SPF and stay out of the sun. Do not take any blood thinners while taking Casodex.
Flomax TAMSULOSIN	0.8 MG TABLET	N/A	ALPHA-BLOCKER THAT RELEASES THE MUSCLES IN THE BLADDER AND PROSTATE AND MAKES IT EASIER TO URINATE.	Do not take with blood thinners. This medication can cause dizziness and headaches. Call for help before getting out of bed. Report any prolonged or painful erections.
TC 99M MEDRONATE	50 mg IV	IV	RADIOACTIVE TRACER THAT ALLOWS PROVIDER TO VIEW AREAS OF ALTERED BONE GROWTH.	Drink plenty of fluids to flush the medication out of your system after administration. Dizziness is common; do not get out of bed without assistance. Immediately report rash or itching to your doctor.