

Due Sunday

IM 7 Math Module

Complete the required math problems and check your answers.

Drop Factor Problems

1. Infuse 1.5 gram of a medication over 3 hours. The drug is supplied as 1.5 gram/250mL. The drip factor is 15. How many gtt/min will you infuse? (Round to the nearest whole number)

$$250\text{mL} \times 15 \div 180 = 21 \text{ gtt/min}$$

2. An order has been written to give 1 gram of a medication over 30 minutes. The drug is supplied as 1 gram/50mL. The gtt factor is 60. How many gtt/min will you infuse?

$$50\text{mL} \times 60 \div 30 = 100 \text{ gtt/min}$$

3. The nurse is to give 500mg IV of a medication over 1 hr. The drug is supplied as 1 gram/250mL. The gtt factor is 15. How many gtt/min will you infuse? (Round to the nearest whole number)

$$250\text{mL} \times 15 \div 60 = 63 \text{ gtt/min}$$

4. The patient is to receive 400 mg IV of a medication over 1 hour. You receive an IV bag from the pharmacy labeled 400 mg in 100 mL D5W. The IV tubing delivers 12 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

$$100\text{mL} \times 12 \div 60 = 20 \text{ gtt/mL}$$

mL/hr Infuse over time

5. The physician writes an order to give 1000mL of intravenous fluid over 8hrs. How many mL/hr will you infuse?

$$1000\text{mL} \div 8\text{hrs} = 125 \text{ mL/hr}$$

6. Infuse 1000 mLs of intravenous fluid over 4 hrs. How many mL/hr will you set on the pump?

$$1000\text{mL} \div 4\text{hrs} = 250 \text{ mL/hr}$$

7. A physician orders 1000 mg of a medication to be given every 6 hours over 1.5 hours. The medication is delivered with 1000 mg in 250 mL. How many mL/hr will you set the pump? (Round to the nearest whole number)

$$250\text{mL} \div 1.5\text{hrs} = 167 \text{ mL/hr}$$

IV Push

8. An order is received for 75mcg IV push of a medication now. The drug is supplied as 100mcg/2mL. How many mL will you give?

$$\frac{75\text{mcg}}{100\text{mcg}} \times 2\text{mL} = 1.5\text{mL}$$

9. The patient is to receive 5mg of a medication IV push. The drug is supplied as 20mg/5mL. How many mL will you give? (Do not round your final answer)

$$\frac{5\text{mg}}{20\text{mg}} \times 5\text{mL} = 1.25\text{mL}$$

10. The order is to give 2mg IV push of a medication now. The drug is supplied as 10mg/1mL. How many mL will you give?

$$\frac{2\text{mg}}{10\text{mg}} \times 1\text{mL} = 0.2\text{mL}$$

IM 7 Math Module

Complete the required math problems and check your answers.

Mcg/kg/min or Mcg/min

$$kg = 79.5$$

11. The patient is receiving an intravenous medication currently infusing at 142 mL/hr. The IV bag of reads 50 mg in 500 mL D5W. The patient weighs 175 lbs. How many mcg/kg/min are infusing? (Round to the nearest tenth)

$$\frac{50\text{mg}}{500\text{mL}} = 0.1 \quad 0.1 \times 1000 = 100 \quad \frac{100 \times 142}{79.5 \times 60} = \frac{14,200}{4,770} = 3 \text{ mcg/kg/min}$$

- 70kg 12. The physician has ordered a medication that states to start at 1 mcg/kg/min and titrate as needed. The IV bag of medication contains 250 mg in 500 mL D5W. The patient weighs 70 kg. How many mL/hr should the IV pump be set at to achieve the starting dose? (Round to the nearest whole number)

$$70\text{kg} \times 1 \text{ mcg/kg/min} \times 60 = \frac{4,200}{500} = 8.4 \text{ mL/hr} \quad \frac{250\text{mg}}{500\text{mL}} = 0.5 \times 1000 = 500 \text{ mcg}$$

13. The patient is currently receiving a medication at 12 mL/hr. The bottle reads 100 mg in 250 mL D5W. How many mcg/min is the patient receiving?

75kg $\frac{100\text{mg}}{250\text{mL}} = 0.4 \quad 0.4 \times 1000 = 400 \quad \frac{400 \times 12}{60} = 80 \text{ mcg/min}$

14. The physician has ordered a medication to start at 2 mcg/kg/min. The patient weighs 165 lbs. The IV bag reads 800 mg in 500 mL D5W. What rate would the nurse set on the infusion pump? (Round to the nearest tenth)

$$\frac{75 \times 2 \times 60}{1600} = \frac{9,000}{1600} = 5.6 \quad \frac{800\text{mg}}{500\text{mL}} = 1.6 \times 1000 = 1600$$

15. The physician has written an order to increase the medication to 4 mcg/kg/min. The IV bag reads 64 mcg/mL. What rate would the nurse set on the IV pump? (Round to the nearest tenth)

$$\frac{4 \times 60}{64} = 3.8 \text{ mL/hr}$$

16. The patient is on a medication drip infusing at 15 mL/hr. The label reads 50 mcg/mL. The patient weighs 65 kg. How many mcg/min is the patient receiving? (Do not round)

$$\frac{50 \times 15}{60} = 12.5$$

Heparin/Insulin or mg/hr

17. The physician orders a heparin infusion at 500 units/hr. The IV bag of medication reads 25,000 units in 250 mL D5W. How many mL/hr should be showing on the IV pump?

$$\frac{25,000 \text{ units}}{250 \text{ mL}} = 100 \quad 500 \div 100 = 5 \text{ mL/hr}$$

18. The patient is on a regular insulin drip infusing at 5 units/hr. The bag is labeled 100 units in 250 mL NS. At what rate should the pump be infusing? Round to the nearest whole number.

$$\frac{100 \text{ units}}{250 \text{ mL}} = 0.4 \quad 5 \div 0.4 = 12.5 \approx 13$$

Burns (Parkland Formula) Do not round weights

19. A 200-pound patient presents to the emergency department with 50% total body surface area (TBSA) burn. How many liters of fluid would be given in the first 24 hours? (Round to the nearest whole number)

$$4 \text{ mL} \times 50 \times 90.9 \div 1000 = 182 \text{ L}$$

20. A 150-pound patient presents to the emergency department with 75% total body surface area (TBSA) burn. How many liters of fluid would be given in the first 24 hours? (Round to the nearest whole number)

$$4 \text{ mL} \times 75 \times 68.2 \div 1000 = 205 \text{ L}$$