

Abriane Trinidad
im7

IM 7 Math Module

Complete the required math problems and check your answers.

Drop Factor Problems

1. Infuse 1.5 gram of a medication over ^{180min} 3 hours. The drug is supplied as 1.5 gram/250mL. The drip factor is 15. How many gtt/min will you infuse? (Round to the nearest whole number)

21 gtt/min

$$\frac{250\text{mL} \times 15\text{gtt}}{180\text{min}} = \frac{3750}{180} = 20.8 = \underline{21\text{gtt/min}}$$

100 gtt/min

2. An order has been written to give 1 gram of a medication over 30 minutes. The drug is supplied as 1 gram/50mL. The gtt factor is 60. How many gtt/min will you infuse?

$$\frac{50 \times 60}{30\text{min}} = \frac{3000}{30} = \underline{100\text{gtt/min}}$$

63 gtt/min

3. The nurse is to give 500mg IV of a medication over ^{60min} 1 hr. The drug is supplied as 1 gram/250mL. The gtt factor is 15. How many gtt/min will you infuse? (Round to the nearest whole number)

$$\frac{250\text{mL} \times 15}{60\text{min}} = \frac{3750}{60} = 62.5 = \underline{63\text{gtt/min}}$$

20 gtt/min

4. The patient is to receive 400 mg IV of a medication over ^{60min} 1 hour. You receive an IV bag from the pharmacy labeled 400 mg in 100 mL D5W. The IV tubing delivers 12 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

$$\frac{100\text{mL} \times 12}{60\text{min}} = \frac{1200}{60} = \underline{20\text{gtt/min}}$$

mL/hr Infuse over time

125 mL/hr

5. The physician writes an order to give 1000mL of intravenous fluid over 8hrs. How many mL/hr will you infuse?

$$\frac{1000\text{mL}}{8\text{hr}} = \underline{125\text{mL/hr}}$$

250 mL/hr

6. Infuse 1000 mLs of intravenous fluid over 4 hrs. How many mL/hr will you set on the pump?

$$\frac{1000\text{mL}}{4\text{hrs}} = \underline{250\text{mL/hr}}$$

IM 7 Math Module

Complete the required math problems and check your answers.

7. A physician orders 1000 mg of a medication to be given every 6 hours over 1.5 hours. The medication is delivered with 1000 mg in 250 mL. How many mL/hr will you set the pump? (Round to the nearest whole number)

107 mL/hr

$$\frac{250 \text{ mL}}{1.5 \text{ hr}} = 166.66 = \underline{107 \text{ mL/hr}}$$

IV Push

8. An order is received for 75 mcg IV push of a medication now. The drug is supplied as 100 mcg/2 mL. How many mL will you give?

1.5 mL

$$\frac{75 \text{ mcg}}{100 \text{ mcg}} \times 2 \text{ mL} = \underline{1.5 \text{ mL}}$$

9. The patient is to receive 5 mg of a medication IV push. The drug is supplied as 20 mg/5 mL. How many mL will you give? (Do not round your final answer)

1.25 mL

$$\frac{5 \text{ mg}}{20 \text{ mg}} \times 5 \text{ mL} = \underline{1.25 \text{ mL}}$$

10. The order is to give 2 mg IV push of a medication now. The drug is supplied as 10 mg/1 mL. How many mL will you give?

0.2 mL

$$\frac{2 \text{ mg}}{10 \text{ mg}} \times 1 \text{ mL} = \underline{0.2 \text{ mL}}$$

Mcg/kg/min or Mcg/min

11. The patient is receiving an intravenous medication currently infusing at 142 mL/hr. The IV bag of reads 50 mg in 500 mL D5W. The patient weighs 175 lbs. How many mcg/kg/min are infusing? (Round to the nearest tenth)

3 mcg/kg/min

$$\frac{\text{conc}}{500 \text{ mg}} \quad \frac{50,000 \text{ mcg}}{500 \text{ mL}} = 100 \text{ mcg/mL} \quad \left| \quad \frac{100 \text{ mcg/mL} \times 142 \text{ mL/hr}}{79.5 \text{ kg} \times 60 \text{ min/hr}} = \frac{14,200}{4770} = 2.97 = \underline{3 \text{ mcg/kg/min}}$$

12. The physician has ordered a medication that states to start at 1 mcg/kg/min and titrate as needed. The IV bag of medication contains 250 mg in 500 mL D5W. The patient weighs 70 kg. How many mL/hr should the IV pump be set at to achieve the starting dose? (Round to the nearest whole number)

8 mL/hr

$$\frac{\text{conc}}{250 \text{ mg}} \quad \frac{250,000}{500 \text{ mL}} = 500 \text{ mcg/mL} \quad \left| \quad \frac{70 \text{ kg} \times 1 \text{ mcg/kg/min} \times 60 \text{ min/hr}}{500 \text{ mcg/mL}} = \frac{4200}{500} = 8.4 = \underline{8 \text{ mL/hr}}$$

13. The patient is currently receiving a medication at 12 mL/hr. The bottle reads 100 mg in 250 mL D5W. How many mcg/min is the patient receiving?

80 mcg/min

$$\frac{\text{conc}}{100 \text{ mg}} \quad \frac{100,000}{250 \text{ mL}} = 400 \text{ mcg/mL} \quad \left| \quad \frac{400 \text{ mcg/mL} \times 12 \text{ mL/hr}}{60 \text{ min/hr}} = \frac{4800}{60} = \underline{80 \text{ mcg/min}}$$

IM 7 Math Module

Complete the required math problems and check your answers.

5.6 mL/hr

14. The physician has ordered a medication to start at 2 mcg/kg/min. The patient weighs 75 kg 165 lbs. The IV bag reads 800 mg in 500 mL D5W. What rate would the nurse set on the infusion pump? (Round to the nearest tenth)

$$\frac{800 \text{ mg}}{500 \text{ mL}} = 1600 \text{ mcg/mL} \quad \left| \quad \frac{75 \text{ kg} \times 2 \text{ mcg/kg/min} \times 60 \text{ min/hr}}{1600 \text{ mcg/mL}} = \frac{9000}{1600} = 5.625 = \underline{5.6 \text{ mL/hr}}$$

3.8 mL/hr

15. The physician has written an order to increase the medication to 4 mcg/kg/min. The IV bag reads 64 mcg/mL. What rate would the nurse set on the IV pump? (Round to the nearest tenth) CONC.

$$\frac{4 \text{ mcg/kg/min} \times 60 \text{ min/hr}}{64 \text{ mcg/mL}} = \frac{240}{64} = 3.75 = \underline{3.8 \text{ mL/hr}}$$

12.5 mcg/min

16. The patient is on a medication drip infusing at 15 mL/hr. The label reads 50 mcg/mL. The patient weighs 65 kg. How many mcg/min is the patient receiving? (Do not round) CONC.

$$\frac{50 \text{ mcg/mL} \times 15 \text{ mL/hr}}{60 \text{ min/hr}} = \frac{750}{60} = \underline{12.5 \text{ mcg/min}}$$

Heparin/Insulin or mg/hr

5 mL/hr

17. The physician orders a heparin infusion at 500 units/hr. The IV bag of medication reads 25,000 units in 250 mL D5W. How many mL/hr should be showing on the IV pump?

$$\frac{500}{25,000} \times 250 \text{ mL} = \underline{5 \text{ mL/hr}}$$

13 mL/hr

18. The patient is on a regular insulin drip infusing at 5 units/hr. The bag is labeled 100 units in 250 mL NS. At what rate should the pump be infusing? Round to the nearest whole number.

$$\frac{5 \text{ u}}{100 \text{ u}} \times 250 \text{ mL} = 12.5 = \underline{13 \text{ mL/hr}}$$

Burns (Parkland Formula) Do not round weights

18L

19. A 200-pound patient presents to the emergency department with 50% total body surface area (TBSA) burn. How many liters of fluid would be given in the first 24 hours? (Round to the nearest whole number)

$$4 \text{ mL} \times 50 \times 90.9 = \frac{18,180}{1000} = 18.18 = \underline{18 \text{ L}}$$

20L

20. A 150-pound patient presents to the emergency department with 75% total body surface area (TBSA) burn. How many liters of fluid would be given in the first 24 hours? (Round to the nearest whole number)

$$4 \text{ mL} \times 75 \times 68.18 = \frac{20,454}{1000} = 20.45 = \underline{20 \text{ L}}$$