

Chantal

## IM 7 Math Module

Complete the required math problems and check your answers.

### Drop Factor Problems

1. Infuse 1.5 gram of a medication over 3 hours. The drug is supplied as 1.5 gram/250mL. The drip factor is 15. How many gtt/min will you infuse? (Round to the nearest whole number)

$$\frac{\text{mL}}{\text{min}} \times \text{gtt} = \underline{21 \text{ gtt/min}}$$

2. An order has been written to give 1 gram of a medication over 30 minutes. The drug is supplied as 1 gram/50mL. The gtt factor is 60. How many gtt/min will you infuse?

$$\frac{50 \text{ mL}}{30 \text{ min}} \times 60 = \underline{100 \text{ gtt/min}}$$

3. The nurse is to give 500mg IV of a medication over 1 hr. The drug is supplied as 1 gram/250mL. The gtt factor is 15. How many gtt/min will you infuse? (Round to the nearest whole number)

$$\frac{\text{mL}}{60 \text{ min}} \times 15 = \frac{250 \text{ mL}}{60 \text{ min}} \times 15 \text{ gtt} = \underline{63 \text{ gtt/min}}$$

4. The patient is to receive 400 mg IV of a medication over 1 hour. You receive an IV bag from the pharmacy labeled 400 mg in 100 mL D5W. The IV tubing delivers 12 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

$$\frac{100 \text{ mL}}{60 \text{ min}} \times 12 = \underline{20 \text{ gtt/min}}$$

mL/hr Infuse over time

5. The physician writes an order to give 1000mL of intravenous fluid over 8hrs. How many mL/hr will you infuse?

$$\frac{1000 \text{ mL}}{8 \text{ hr}} = \underline{125 \text{ mL/hr}}$$

6. Infuse 1000 mLs of intravenous fluid over 4 hrs. How many mL/hr will you set on the pump?

$$\frac{1000 \text{ mL}}{4 \text{ hr}} = \underline{250 \text{ mL/hr}}$$

7. A physician orders 1000 mg of a medication to be given every 6 hours over 1.5 hours. The medication is delivered with 1000 mg in 250 mL. How many mL/hr will you set the pump? (Round to the nearest whole number)

$$\frac{250 \text{ mL}}{1.5 \text{ hrs}} = \underline{167 \text{ mL/hr}}$$

IV Push

8. An order is received for 75mcg IV push of a medication now. The drug is supplied as 100mcg/2mL. How many mL will you give?

$$\frac{75 \text{ mcg}}{100 \text{ mcg}} \times 2 \text{ mL} = \underline{1.5 \text{ mL}}$$

9. The patient is to receive 5mg of a medication IV push. The drug is supplied as 20mg/5mL. How many mL will you give? (Do not round your final answer)

$$\frac{5 \text{ mg}}{20 \text{ mg}} \times 5 \text{ mL} = \underline{1.25 \text{ mL}}$$

10. The order is to give 2mg IV push of a medication now. The drug is supplied as 10mg/1mL. How many mL will you give?

$$\frac{2 \text{ mg}}{10 \text{ mg}} \times 1 \text{ mL} = \underline{0.2 \text{ mL}}$$

## IM 7 Math Module

Complete the required math problems and check your answers.

Mcg/kg/min or Mcg/min

11. The patient is receiving an intravenous medication currently infusing at 142 mL/hr. The IV bag of reads 50 mg in 500 mL D5W. The patient weighs 175 lbs. How many

mcg/kg/min are infusing? (Round to the nearest tenth)  
 $\frac{50 \text{ mg}}{500 \text{ mL}} = 0.1 \text{ mg/mL} = 100 \text{ mcg/mL}$   
 $\frac{100 \text{ mcg} \times 142 \text{ mL/hr}}{60 \text{ min} \times 19.5 \text{ kg}} = \frac{14,200}{1,170} = 3 \text{ mcg/kg/min}$  or 3.0

12. The physician has ordered a medication that states to start at 1 mcg/kg/min and titrate

as needed. The IV bag of medication contains 250 mg in 500 mL D5W. The patient weighs 70 kg. How many mL/hr should the IV pump be set at to achieve the starting dose? (Round to the nearest whole number)

$\frac{250 \text{ mg}}{500 \text{ mL}} = 0.5 \text{ mg/mL} = 500 \text{ mcg}$   
 $\frac{1 \text{ mcg/kg/min} \times 70 \text{ kg} \times 60 \text{ min}}{500 \text{ mcg}} = 8 \text{ mL/hr}$

13. The patient is currently receiving a medication at 12 mL/hr. The bottle reads 100 mg in 250 mL D5W. How many mcg/min is the patient receiving?

$\frac{100 \text{ mg}}{250 \text{ mL}} = 0.4 \text{ mg/mL} = 400 \text{ mcg/mL}$   
 $\frac{400 \text{ mcg/mL} \times 12 \text{ mL/hr}}{60 \text{ min/hr}} = 80 \text{ mcg/min}$

14. The physician has ordered a medication to start at 2 mcg/kg/min. The patient weighs

165 lbs. The IV bag reads 800 mg in 500 mL D5W. What rate would the nurse set on the infusion pump? (Round to the nearest tenth)

$\frac{800 \text{ mg}}{500 \text{ mL}} = 1.6 \text{ mg/mL} = 1600 \text{ mcg/mL}$   
 $\frac{2 \text{ mcg/kg/min} \times 75 \text{ kg} \times 60 \text{ min}}{1600 \text{ mcg/mL}} = 5.6 \text{ mL/hr}$

15. The physician has written an order to increase the medication to 4 mcg/kg/min. The IV bag reads 64 mcg/mL. What rate would the nurse set on the IV pump? (Round to the nearest tenth)

$\frac{4 \text{ mcg/kg/min} \times 60 \text{ min}}{64 \text{ mcg/mL}} = 3.8 \text{ mL/hr}$

16. The patient is on a medication drip infusing at 15 mL/hr. The label reads 50 mcg/mL. The patient weighs 65 kg. How many mcg/min is the patient receiving? (Do not round)

$\frac{50 \text{ mcg/mL} \times 15 \text{ mL/hr}}{60 \text{ min}} = \frac{750}{60} = 12.5 \text{ mcg/min}$

Heparin/Insulin or mg/hr

17. The physician orders a heparin infusion at 500 units/hr. The IV bag of medication reads 25,000 units in 250 mL D5W. How many mL/hr should be showing on the IV pump?

$\frac{500 \text{ units}}{25,000 \text{ units}} \times 250 \text{ mL} = 5 \text{ mL/hr}$

18. The patient is on a regular insulin drip infusing at 5 units/hr. The bag is labeled 100 units in 250 mL NS. At what rate should the pump be infusing? Round to the nearest whole number.

$\frac{5 \text{ units}}{100 \text{ units}} \times 250 \text{ mL} = 12.5 \text{ mL/hr}$

Burns (Parkland Formula) Do not round weights

19. A 200-pound patient presents to the emergency department with 50% total body surface area (TBSA) burn. How many liters of fluid would be given in the first 24 hours? (Round to the nearest whole number)

$4 \text{ mL} \times 50\% \times 90.9 \text{ kg} = 18,180 \text{ mL} = 18.18 \text{ L} = 18 \text{ L}$

20. A 150-pound patient presents to the emergency department with 75% total body surface area (TBSA) burn. How many liters of fluid would be given in the first 24 hours? (Round to the nearest whole number)

$4 \text{ mL} \times 75\% \times 68.18 \text{ kg} = 20,454 \text{ mL} = 20.454 \text{ L} = 20 \text{ L}$

$4 \text{ mL} \times 75\% \times 68.18 \text{ kg} = 20,454 \text{ mL} = 20.454 \text{ L} = 20 \text{ L}$