

Student Name: Cynthia A. Cerda

Unit: Pedi 3

Pt. Initials: CCS

Date: 1-29-2025

Pediatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Allergies: NKA

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Therapeutic Range?	IVP – List diluent solution, volume, and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				Is med in therapeutic range? If not, why?			
Amlodipine (Norvasc)	Calcium Channel Blocker	↓ BP (HTN redm)	5mg PO Daily	2.5mg IV daily Yes Max 10		Swelling of feet/ankles Redness of face, neck, arms	1. Get up slowly because this med can make you dizzy/light headed due to changes in blood pressure 2. Use the call light so I can be with you to make sure you know how drugs will affect you (normal falls) 3. May make you drowsy because of effect on BP 4. Let us know if you have swelling or feel faint ASAP
Ceftriaxone (Rocephin)	Antibiotic	Fights to kill bacteria	2000mg in sterile H <sub>2</sub> O 20mL syng q24hrs	50-75mg/kg/day in divided doses q 12-24hrs Max daily 2000mg/day Yes	IVP over 3-5 min Do NOT use calcu casts IVs (LR) + TPNA Yes (2000mg in sterile water 20mL syng (100mg/mL)	Diarrhea Pneumonia Black/tarry stools	1. Call dr. if you have watery/bloody diarrhea anytime if med. This may cause infectious diarrhea which would require treatment 2. Take all of this medicine unless your care team tells you otherwise keep taking it even if you feel better 3. If you experience confusion, increasing loss of appetite, thinking clearly, notify physician 4. If you feel short of breath let us know
Famotidine (Pepcid)	Acid Reducer	Reduce Acid	20mg tab BID	10-20mg q 12hrs Max 2 tabs/day Yes		Headache/Dizziness Constipation or Diarrhea	1. Contact Dr. if you feel dizzy/like you are going to pass out 2. Let me know if you feel a racing heartbeat 3. Do not chew this med 4. Call Dr. if you have a peeling rash/blistered skin, wheezing/tightness of chest
Methylprednisolone Sodium Succinate (Solu Medrol)	Adrenal Corticosteroid Anti-inflammatory	Anti-inflammatory	40mg/mL 1mg 30mg q 12hrs	0.11-1.6mg/kg/day in 3-4 doses Yes	Mix 1mL of sterile H <sub>2</sub> O to make 40 mg/mL over 5 min	Fluid retention Wt gain ↑ BP Potassium loss	1. Take with/after food to minimize stomach upset 2. Do not stop this med abruptly due to risk of adrenal insufficiency syndrome - may ↑ blood glucose 3. May cause mood swings + insomnia because this med affects cortisol which plays a critical part in our sleep/wake cycle 4. Cortisol in this med can raise the stress hormone - it controls hunger (part of the brain) 5. In stool - indicates diarrhea/infection 6. ↑ Gas 7. Stomach pain that gets worse, whitish or yellowish stool may indicate liver issues 8. Swelling/persistent diarrhea may indicate inflammation
Pantoprazole (Protonix)	Gastric Secretion Inhibitor Proton Pump Inhibitor	↓ GI reflux ↓ acid in stomach	40mg in NaCl 0.9% inj in 10mL syng q 24hrs	2.5mg/kg/day 1.2mg/kg/dose in 30 min before meal Yes	IVP over 2 min 40mg in NaCl (0.9%) inj 10mL syng.	Headache Nausea Diarrhea	1. Call dr. if you have persistent diarrhea/blood in stool - indicates diarrhea/infection 2. ↑ Gas 3. Stomach pain that gets worse, whitish or yellowish stool may indicate liver issues 4. Swelling/persistent diarrhea may indicate inflammation